

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, C07K 14/00	A1	(11) International Publication Number: WO 98/15648 (43) International Publication Date: 16 April 1998 (16.04.98)									
(21) International Application Number: PCT/DK97/00425 (22) International Filing Date: 3 October 1997 (03.10.97) (30) Priority Data: <table border="0"> <tr> <td>1096/96</td> <td>4 October 1996 (04.10.96)</td> <td>DK</td> </tr> <tr> <td>1156/96</td> <td>18 October 1996 (18.10.96)</td> <td>DK</td> </tr> <tr> <td>0512/97</td> <td>5 May 1997 (05.05.97)</td> <td>DK</td> </tr> </table> (71) Applicant: DAKO A/S [DK/DK]; Produktionsvej 42, DK-2600 Glostrup (DK). (72) Inventors: STENDER, Henrik; Fasanhaven 5, DK-2820 Gentofte (DK). LUND, Kaare; A.D. Jørgensensvej 19 3 tv, DK-2000 Frederiksberg (DK). MOLLERUP, Tina, Andersen; Lejrevej 14, Allerslev, DK-4320 Lejre (DK). (74) Agent: FINK, Kirsten; Dako a/s, Produktionsvej 42, DK-2600 Glostrup (DK).		1096/96	4 October 1996 (04.10.96)	DK	1156/96	18 October 1996 (18.10.96)	DK	0512/97	5 May 1997 (05.05.97)	DK	(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
1096/96	4 October 1996 (04.10.96)	DK									
1156/96	18 October 1996 (18.10.96)	DK									
0512/97	5 May 1997 (05.05.97)	DK									
(54) Title: NOVEL PROBES FOR THE DETECTION OF MYCOBACTERIA (57) Abstract <p>Novel hybridisation assay probes and mixtures of such probes for detecting a target sequence of one or more mycobacteria optionally present in a sample. The probes may suitably be directed to target sequences of mycobacterial rDNA, precursor rRNA, or rRNA, said probes being capable of forming detectable hybrids. The probes are in particular directed to mycobacterial rDNA, to precursor rRNA, or to 23S, 16S or 5S rRNA. The probes are useful for detecting the organisms in test samples such as sputum, laryngeal swabs, gastric lavage, bronchial washings, biopsies, aspirates, expectorates, body fluids (spinal, pleural, pericardial, synovial, blood, pus, bone marrow), urine, tissue sections as well as food samples, soil, air and water samples, and cultures thereof.</p>											

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NOVEL PROBES FOR THE DETECTION OF MYCOBACTERIA

The present invention relates to novel probes and to mixtures of such probes, in addition to the design, construction and use of such novel probes or mixtures thereof for detecting a target sequence of one or more mycobacteria, which probes are capable of detecting such organism(s) optionally present in a test sample, e.g. sputum, laryngeal swabs, gastric lavage, bronchial washings, biopsies, aspirates, expectorates, body fluids (spinal, pleural, pericardial, synovial, blood, pus, bone marrow), urine, tissue sections as well as food samples, soil, air and water samples and cultures thereof. The invention relates in particular to novel probes and mixtures thereof for detecting the presence of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC) and for detecting the presence of one or more mycobacteria other than mycobacteria of the Mycobacterium tuberculosis Complex (MOTT). The invention further relates to diagnostic kits comprising one or more of such probes. The probes of the present invention are surprisingly able to penetrate the cell wall of the mycobacteria, thus making possible the development of fast and easy-performed in situ protocols.

BACKGROUND OF THE INVENTION

Tuberculosis is a very life-threatening and highly epidemic disease which is caused by infection with Mycobacterium tuberculosis. Tuberculosis is presently the predominant infectious cause of morbidity and mortality world-wide, and is estimated to kill about three million people annually. WHO estimates that the annual number of new cases of tuberculosis will increase from 7.5 million in 1990 to 10.2 million in 2000, an escalation that will result in approximately 90 million new cases during this decade. It is furthermore estimated that 30 million people will die from tuberculosis during the 1990s, which equals one quarter of preventable deaths among adults.

The prevalence of tuberculosis has been very high in the poorer parts of the world such as Asia, Africa and South-America, but in recent years an increase has also been observed in industrialised countries. This appears to be due to an interaction of various factors including i.a. patterns of migration, poorly organised tuberculosis programmes and nutrition problems. Furthermore, a serious threat will arise from the emergence of new strains that are drug resistant or worse, multi-drug resistant.

Mycobacteria are often divided into tuberculous mycobacteria, i.e. mycobacteria of the Mycobacterium tuberculosis Complex (MTC), and non-tuberculous mycobacteria, i.e. mycobacteria other than those of the Mycobacterium tuberculosis Complex (MOTT). The MTC

group comprises apart from *M. tuberculosis*, *M. bovis*, *M. africanum* and *M. microti*.

Mycobacteria of the MOTT group are not normally pathogenic to healthy individuals but may cause disease in immunocompromised individuals, e.g. individuals infected with HIV. Clinical relevant mycobacteria of the MOTT group are in particular *M. avium*, *M. intracellulare*, *M.*

5 *kansasii* and *M. gordonae*, but also *M. scrofulaceum*, *M. xenopi* and *M. fortuitum*.

M. avium and *M. intracellulare* together with *M. paratuberculosis* and *M. lepraemurium* constitute the Mycobacterium avium Complex (MAC). Extended with *M. scrofulaceum*, the group is named Mycobacterium avium -intracellulare -scrofulaceum Complex (MAIS).

10

It is well-known that treatment of mycobacterial infections with antibiotics may lead to the emergence of drug resistant strains. Many antibiotic drugs exert their effects by interfering with protein synthesis or with transcription. Studies of the molecular mechanisms underlying certain antibiotic resistance phenotypes in clinical mycobacterium isolates have revealed

15 mutations in rRNA genes. The development of resistance because of mutation(s) located in the rRNA gene is likely to occur since slow-growing mycobacteria have only a single rRNA operon. All mycobacteria populations comprise a minority of drug resistant mutants that have arisen by spontaneous mutation. These mutated mycobacteria do normally not survive particularly well, but, when single-drug therapy is offered as treatment, the drug susceptible

20 bacteria are killed, and only the resistant mutants will survive and multiply, and, thus at some point, constitute the majority of the mycobacterial population. The selection of drug resistant bacteria due to inadequate drug therapy leads to a state of so-called "acquired drug-resistance". In contrast, "primary drug-resistance" is used to characterise a situation where drug-resistant mycobacteria can be isolated from a patient who has never been treated for

25 mycobacterial infection, and has become infected with drug-resistant mycobacteria from an individual suffering from infection with an acquired drug resistant bacterium.

Today, drug-resistance is determined primarily phenotypically by culturing clinical samples, in which presence of mycobacteria have been demonstrated, in the presence of the individual

30 drugs. This is unfortunately a very slow and time-consuming procedure as the result of the drug-resistance studies depends on the growth rate of the mycobacteria, which are well-known to be slow. Thus, the result is not available until after several weeks.

Although the incidence of drug-resistance is, at least not yet, very common, it is nevertheless

35 very important that resistant strains are identified and eradicated. Therefore, it is of major importance to find a reliable and rapidly performed method of diagnosing drug-resistance.

Presently, the detection of mycobacteria by microscopy is the most prevalent method for

diagnosis. The sample (e.g. an expectorate) is stained for the presence of acid-fast bacilli using e.g. Ziehl-Neelsen staining. However, staining for acid-fast bacilli does not provide the necessary information about the type of infection, only whether acid fast bacilli are present in the sample, and this is in itself not sufficient information for establishing a diagnosis. Samples
5 positive for acid fast bacilli, may subsequently be cultured in order to be able to perform species identification.

Since Ziehl-Neelsen staining cannot be used to determine whether the infection is caused by mycobacteria of the MTC group or mycobacteria other than mycobacteria of the MTC group, a
10 positive staining frequently leads to very costly isolation of all the patients with suspected M. tuberculosis infection as well as treatment with medicaments to which the patient may not even respond.

Since the sensitivity of acid fast staining is only approximately 10^4 - 10^5 per ml smear negative
15 samples should also be cultured as culture-based tests are sensitive, and as it may be possible to detect 10-100 organisms per sample, but the result is not available before up to 8 weeks of culturing. Likewise, information about drug susceptibility is not available until after 1-3 weeks of further testing.

20 Different solid or liquid media (Loewenstein Jensen slants and Dubos broth) have traditionally been used for culturing of mycobacteria-containing samples. Newer media include ESP Myco Culture System (Difco), MB/BacT (Organon Teknika), BacTec (Becton Dickinson) and MGIT (Becton Dickinson). These test media are based on colourmetric or fluorometric detection of carbon dioxide or oxygen produced by mycobacterial metabolism, and adapted to automated
25 systems for large scale testing.

Species identification is presently carried out following culturing using traditional biochemical methods or probe hybridisation assays (e.g. AccuProbe by Gen-Probe Inc., USA). There is, therefore, an increasing need for means allowing a more rapid distinction between
30 mycobacteria of the MTC group and mycobacteria other than those of the MTC group, and for further species identification of those especially mycobacteria other than those of the MTC group.

A number of new attempts to replace the culture-based methods relies on molecular
35 amplification technology. Several methods have emerged, among them the polymerase chain reaction (PCR), the ligase chain reaction and transcription mediated amplification. The basic principle of amplification methods is that a specific nucleic acid sequence of the mycobacteria is amplified to increase the copy number of the specific sequence to a level where the

amplicon may be detectable. In principle, the methods offers the possibility of detecting only one target sequence, thus, in principle, making detection of mycobacteria present at low levels possible. However, it has become clear that the target amplification methods cannot replace culture-based methods as only samples which are positive by staining for acid fast bacilli (AFB) give a satisfactory sensitivity. Furthermore, specific problems exist for each method. The PCR method may give false negative results due to the presence of inhibitors such as haemoglobin. Another problem arises from cross-contamination of negative specimens and/or reagents with amplified nucleic acid present in the laboratory environment leading to false positive results. A disadvantage is that costly reagents are needed for performing these tests. Furthermore, specialised instrumentation is required, making these tests mainly useful in large specialised laboratories, and generally not applicable in smaller clinical laboratories.

Nucleic acid probes for detecting rRNA of mycobacteria have been described in for example US 5 547 842, EP-A 0 572 120 and US 5 422 242.

Considering the perspective and impact the disease has, the development of rapid and preferably easy-performed and further economic feasible diagnostic detection tests are of utmost importance and would be a very valuable tool in the fight against the spread of tuberculosis.

Peptide nucleic acids are pseudo-peptides with DNA-binding capability. The compounds were first reported in the early nineties in connection with a series of attempts to design nucleotide analogues capable of hybridising, in a sequence-specific fashion, to DNA and RNA, cf. WO 92/20702.

Hybridisation of peptide nucleic acid probes to DNA and to RNA has been shown to obey the Watson-Crick base pairing rules, and peptide nucleic acid probes have been found to hybridise to a DNA or a RNA target with higher affinity and specificity than the nucleic acid counterparts. These properties are ascribed to the uncharged, as opposed to the charged, structure of the peptide nucleic acid and DNA or RNA backbones, respectively, and to the high conformational flexibility of the peptide nucleic acid molecules. These features - together with the documented stability of peptide nucleic acid towards a variety of naturally occurring nucleases and proteases that usually degrade DNA, RNA or proteins - invite for use of peptide nucleic acid probes as antisense therapeutic agents and opens potentially important applications in diagnostics.

SUMMARY OF THE INVENTION

The present invention relates to novel peptide nucleic acid probes and to mixtures of such probes for detecting a target sequence of one or more mycobacteria optionally present in a sample. In accordance with claim 1, the probes are directed to target sequences of mycobacterial rRNA, genomic sequences corresponding to said rRNA (rDNA) and precursor
5 rRNA. rRNA is present in a high number of copies in each cell, and is hence a well suited target. The probes are, as defined in claim 2, suitably directed to target sequences of mycobacterial rDNA, precursor rRNA, or 23S, 16S or 5S rRNA.

Thus, in a first aspect, the invention features a hybridisation assay probe and a mixture of
10 such probes for detecting a target sequence of one or more mycobacteria in accordance with claim 1 and 2. Under appropriate stringency conditions, Such probes should not to any significant degree cross-react with ribosomal nucleic acid from other not relevant organisms, present in the test sample, in particular other mycobacteria. Cross-reactivity to organisms that are unlikely to be present in the sample may not be of importance. In in situ assays implying
15 examination by microscopy, it is further possible to distinguish mycobacteria from other bacteria based on the morphology of these bacilli.

The invention also relates to peptide nucleic acid probes in accordance with claim 3 for obtaining a target sequence and in accordance with claim 4 for obtaining a probe.
20

In another aspect, the invention relates to novel peptide nucleic acid probes for detecting a target sequence of one or more mycobacteria of the MTC group, and one or more mycobacteria other than mycobacteria of the MTC group, which probes comprise from 6 to 30 polymerised peptide nucleic acid moieties (claim 5). Suitable probes of formula (I) are claimed
25 in claim 6.

Claims 7 to 10 and 15 to 24 relate to probes or mixtures of such probes for detecting a target sequence of one or more mycobacteria of the MTC group. Claims 11 to 13 and 15 to 24 relate to probes or mixtures of such probes for detecting a target sequence of one or more
30 mycobacteria other than mycobacteria of the MTC group (MOTT group). Claim 14 relates specifically to probes for detecting drug resistant mycobacteria. Claims 25 to 27 relate to the use of such probes or mixtures thereof.

In accordance with claims 28 to 34, the present invention also relates to a method for
35 detecting the presence of mycobacteria.

In yet another aspect, the present invention relates to a kit (claim 35 and 36) comprising at least one peptide nucleic acid probe as defined in claims 1 to 24.

Mycobacteria are characterised by a complex cell wall which contains mycolic acids, complex waxes and unique glycolipids. It is generally recognised by those skilled in the art that this wall provides mycobacteria with extreme resistance to chemical and physical stress as compared to other bacteria, and, accordingly, makes them very difficult to penetrate and lyse. The low permeability of the cell wall is considered to be the main reason for the fact that only very few drugs are effective in the treatment of tuberculosis and other mycobacterial infections. As described in US 5 582 985, the wall appears further to prevent penetration by nucleic acid probes. Even with short probes (shorter than 30 nucleic acids), specific staining is low or often non-existent. Protocols that allow DNA probes to be used for in situ hybridisation to mycobacterial species are described in US 5 582 985. However, these protocols require dewaxing of the mycobacterial cell wall with xylene and further enzymatic treatment prior to the hybridisation step in order to make the mycobacterial cell wall permeable to the DNA probes.

The problems set forth above have surprisingly been completely solved by the use of peptide nucleic acid probes. It has, surprisingly, been found that the peptide nucleic acid probes are able to penetrate the cell wall of the mycobacteria, and furthermore that this is taking place rapidly. The person skilled in the art would arrive at the conviction that it would be necessary to heavily treat the mycobacteria before hybridisation is carried out. Thus, based on the available prior art, there is a strong prejudice against carrying out hybridisation without prior destruction of the mycobacterial cell wall. The inventors have shown that this is indeed and unexpectedly possible. It has been demonstrated that the probes of the present invention are able to hybridise to mycobacterial precursor rRNA and rRNA without harsh treatment of the mycobacterial cells, thus avoiding a risk of interfering with the morphology of the cells. Using the present probes, specific and easy detection and, subsequently, diagnosis of tuberculosis and other mycobacterial infections are thus possible.

BRIEF DESCRIPTION OF THE FIGURES

Alignments of rDNA sequences of *M. tuberculosis* (as a representative of the MTC group) and important closely related species thereto, including *M. avium* (as a representative of the mycobacteria other than those of the MTC group) and important closely related species thereto for the 23S, 16S and/or 5S rRNA genes have been made (Figures 1A-1J, 2A-2D, 3, 4A-4L and 5A-B). The alignment for *M. bovis* and *M. intracellulare* are partly based on public available sequences and partly on sequences obtained by sequencing performed at DAKO A/S.

Alignment for the MTC group (23S rDNA)

Figures 1A-1J show alignments of 23S rDNA sequences of *M. tuberculosis* (GenBank entry GB:MTCY130, accession number Z73902), *M. avium* (GenBank entry GB:MA23SRNA, accession number X74494), *M. paratuberculosis* (GenBank entry GB:MPARRNA, accession number X74495), *M. phlei* (GenBank entry GB:MP23SRNA, accession number X74493), *M. leprae* (GenBank entry GB:ML5S23S, accession number X56657), *M. gastri* (GenBank entry GB:MG23SRRNA, accession number Z17211), *M. kansasii* (GenBank entry GB:MK23SRRNA, accession number Z17212), and *M. smegmatis* (GB:MS16S23S5, accession number Y08453). Preferred peptide nucleic acid probes should enclose at least one nucleobase complementary to a nucleobase of *M. tuberculosis* 23S rRNA within positions 149-158, 220-221, 328-361, 453-455, 490-501, 637-660, 706-712, 762-789, 989, 1068-1072, 1148, 1311-1329, 1361-1364, 1418, 1563-1570, 1627-1638, 1675-1677, 1718, 1734-1740, 1967-1976, 2403-2420, 2457-2488, 2952-2956, 2966-2969, 3000-3003, and 3097-3106 of the alignment (indicated by heavy frames). Differences between the sequences of *M. avium*, *M. phlei*, *M. leprae*, *M. paratuberculosis*, *M. gastri* and *M. kansasii* and that of *M. tuberculosis* in the alignment are indicated by light frames.

Alignment for the MTC group (16S rDNA)

Figures 2A-2D show alignments of 16S rDNA sequences of *M. tuberculosis* (GenBank entry GB:MTU16SRN, accession number X52917), *M. bovis* (GenBank entry GB:MSGTGDA, accession number M20940), *M. avium* (GenBank entry GB:MSGRRDA, accession number M61673), *M. intracellulare* (GenBank entry GB:MIN16SRN, accession number X52927), *M. paratuberculosis* (GenBank entry GB:MSGRRDH, accession number M61680), *M. scrofulaceum* (GenBank entry GB:MSC16SRN, accession number X52924), *M. leprae* (GenBank entry GB:MLEP16S1, accession number X55587), *M. kansasii* (GenBank entry GB:MKRRN16, accession number X15916), *M. gastri* (GenBank entry GB:MGA16SRN, accession number X52919), *M. gordonae* (GenBank entry GB:MSGRR16SI, accession number M29563) and *M. marinum* (GenBank entry GB:MMA16SRN, accession number X52920). Preferred peptide nucleic acid probes should enclose at least one nucleobase complementary to a nucleobase of *M. tuberculosis* 16S rRNA within positions 76-79, 98-101, 135-136, 194-201, 222-229, 242, 474, 1136-1145, 1271-1272, 1287-1292, 1313, and 1334 of the alignment (indicated by heavy frames). Differences between the sequences of *M. bovis*, *M. avium*, *M. intracellulare*, *M. paratuberculosis*, *M. scrofulaceum*, *M. leprae*, *M. kansasii*, *M. gastri*, *M. gordonae* and *M. marinum*, and that of *M. tuberculosis* in the alignment are indicated by light frames.

Alignment for the MTC group (5S rDNA)

Figure 3 shows alignments of 5S rDNA sequences of *M. tuberculosis* (GenBank entry

GB:MTDNA16S, accession number x75601), *M. bovis* (GenBank entry GB:MBRRN5S, accession number X05526), *M. phlei* (GenBank entry GB:MP5SRRNA, accession number X55259), *M. leprae* (GenBank entry GB:ML5S23S, accession number X56657), and *M. smegmatis* (GenBank entry GB:MS16S23S5, accession number Y08453). Preferred peptide
 5 nucleic acid probes should enclose at least one nucleobase complementary to a nucleobase of *M. tuberculosis* 5S rRNA within positions 86-90 of the alignment (indicated by heavy frame). Differences between the sequences of *M. bovis*, *M. phlei*, *M. leprae*, *M. smegmatis* and *M. luteus* and that of *M. tuberculosis* in the alignment are indicated by light frames.

10 *Alignment for Mycobacteria other than those of the MTC group (23S rDNA)*

Figures 4A-4L show alignments of 23S rDNA sequences of *M. avium* (GenBank entry GB:MA23SRNA, accession number X74494), *M. paratuberculosis* (GenBank entry GB:MPARRNA, accession number X74495), *M. tuberculosis* (GenBank entry GB:MTCY130, accession number Z73902), *M. phlei* (GenBank entry GB:MP23SRNA, accession number
 15 X74493), *M. leprae* (GenBank entry GB:ML5S23S, accession number X56657), *M. gastris* (GenBank entry GB:MG23SRRNA, accession number Z17211), *M. kansasii* (GenBank entry GB:MK23SRRNA, accession number Z17212), and *M. smegmatis* (GB:MS16S23S5, accession number Y08453). Preferred peptide nucleic acid probes should enclose at least one nucleobase complementary to a nucleobase of *M. avium* 23S rRNA within positions 99-101,
 20 183, 261-271, 281-284, 290-293, 327-335, 343-357, 400-405, 453-462, 587-599, 637-660, 704-712, 763-789, 1060-1074, 1177-1185, 1259-1265, 1311-1327, 1345-1348, 1361-1364, 1556-1570, 1608-1613, 1626-1638, 1651-1659, 1675-1677, 1734-1741, 1847-1853, 1967-1976, 2006-2010, 2025-2027, 2131-2232, 2252-2255, 2396-2405, 2416-2420, 2474-2478, 2687, 2719, 2809, 3062-3068, and 3097-3106 of the alignment (indicated by heavy frames).
 25 Differences between the sequences of *M. paratuberculosis*, *M. tuberculosis*, *M. phlei*, *M. leprae*, *M. gastris*, *M. kansasii*, and *M. smegmatis* and that of *M. avium* in the alignment are indicated by light frames.

Alignment for Mycobacteria other than those of the MTC group (16S rDNA)

30 Figures 5A-5B show alignments of 16S rDNA sequences of *M. avium* (GenBank entry GB:MSGRRDA, accession number M61673), *M. intracellulare* (GenBank entry GB:MIN16SRN, accession number X52927), *M. paratuberculosis* (GenBank entry GB:MSGRRDH, accession number M61680), *M. scrofulaceum* (GenBank entry GB:MSC16SRN, accession number X52924), *M. tuberculosis* (GenBank entry GB:MTU16SRN, accession number X52917), *M. bovis* (GenBank entry GB:MSGTGDA, accession number
 35 M20940), *M. leprae* (GenBank entry GB:MLEP16S1, accession number X55587), *M. kansasii* (GenBank entry GB:MKRRN16, accession number X15916), and *M. gastris* (GenBank entry GB:MGA16SRN, accession number X52919), *M. gordonae* (GenBank entry GB:MSGRR16S1,

- accession number M29563), and *M. marinum* (GenBank entry GB:MMA16SRN, accession number X52920). Preferred peptide nucleic acid probes should enclose at least one nucleobase complementary to a nucleobase of *M. avium* 16S rRNA within positions 135-136, 472-475, 1136-1144, 1287-1292, 1313, and 1334 of the alignment (indicated by heavy frames). Differences between the sequences of *M. intracellulare*, *M. paratuberculosis*, *M. scrofulaceum*, *M. tuberculosis*, *M. bovis*, *M. leprae*, *M. kansasii*, and *M. gastri* and that of *M. avium* in the alignment are indicated by light frames.

Drug-resistance

- Figure 6 shows a partial *M. avium* 23S rDNA sequence including positions 2550 to 2589 of GenBank entry X74494. Bases in positions where deviations from the wild-type sequence have been correlated with macrolide-resistance are framed. Positions 2568 and 2569 in the figure correspond to positions 2058 and 2059, respectively, of *E. coli* 23S rRNA.
- Figure 7 shows a partial *M. tuberculosis* 16S rDNA sequence including positions 441 to 491 and 843 to 883 of GenBank entry X52917. Bases in positions where deviations from the wild-type sequence have been correlated with resistance to streptomycin are framed. Positions 452, 473, 474, 477, 865, and 866 in the figure correspond to positions 501, 522, 523, 526, 912, and 913, respectively, of *E. coli* 16S rRNA.

20

SPECIFIC DESCRIPTION

- The present invention provides novel probes for use in rapid and specific, sensitive hybridisation based assays for detecting a target sequence of one or more mycobacteria, which target sequence is located in the mycobacterial rDNA, precursor rRNA, or in the 23S, 16S or 5S rRNA. The probes to be used in accordance with the present invention are peptide nucleic acid probes. Peptide nucleic acids are non-naturally occurring polyamides or polythioamides which can bind to nucleic acids (DNA and RNA). Such compounds are described in e.g. WO 92/20702.

30

- We have identified suitable variable regions of the target nucleic acid by comparative analysis of generally available rDNA sequences and sequences obtained by sequencing as described above. Computers and computer programs, which have been used for the purposes disclosed herein, are commercially available. From such alignments, possibly suitable probes can be identified. The alignments are thus a useful guideline for designing probes with desired characteristics.

35

When designing the probes, due regard should be taken to the assay conditions under which

the probes are to be used. Stringency is chosen so as to maximise the difference in stability between the hybrid formed with the target nucleic acid and that formed with the non-target nucleic acid. It will typically be necessary to choose high stringency conditions for probes where the specificity depends on only one mismatch to non-target sequences. The more
5 mismatches to non-target sequences, the less demand for high stringency conditions.

Furthermore, probes should be designed so as to minimise the stability of probe-non-target nucleic acid hybrids. This may be accomplished by minimising the degree of complementarity to non-target nucleic acid, i.e. by designing the probe to span as many destabilising
10 mismatches as possible, and/or to include as many additions/deletions relative to the target sequence as possible. Whether a probe is useful for detecting a particular mycobacterial species depends to some degree on the difference between the thermal stability of probe-target hybrids and probe:non-target hybrids. For rRNA targets, however, the secondary structure of the region of the rRNA molecule in which the target sequence is located may also
15 be of importance. The secondary structure of a probe should also be taken into consideration. Probes should be designed so as to minimise their proclivity to form hairpins, self-dimers, and pair-dimers if a mixture of two or more probes is used.

Mismatching bases in hybrids formed between peptide nucleic acid probes and nucleic acids
20 result in a higher thermal instability than mismatching bases in nucleic acid duplexes of the same sequences. Thus, the peptide nucleic acid probes exhibit a greater specificity for a given target nucleic acid sequence than a traditional nucleic acid probe, which is seen as a greater difference in T_m values for probe-target hybrids and probe-non-target hybrids. The sensitivity and specificity of a peptide nucleic acid probe will also depend on the hybridisation conditions
25 used.

The primary concern regarding the length of the peptide nucleic acid probes is the warranted specificity, i.e. which length provides sufficient specificity for a particular application. The optimal length of a peptide nucleic acid probe comprising a particular site with differences in
30 base composition, e.g. among selected regions of mycobacterial rRNA, is a compromise between the general pattern that longer probes ensure specificity and shorter probes ensure that the destabilising differences in base composition constitute a greater portion of the probe. Also, due regard must be paid to the conditions under which the probes are to be used.

35 Peptide nucleic acid sequences are written from the N-terminal end of the sequence towards the C-terminal end. A free (unsubstituted) N-terminal end or an N-terminal end terminating with an amino acid is indicated as H, and a free C-terminal end is indicated as NH_2 (a carboxamide group). Peptide nucleic acids are capable of hybridising to nucleic acid

sequences in two orientations, namely in antiparallel orientation and in parallel orientation. The peptide nucleic acid is said to hybridise in the antiparallel orientation when the N-terminal end of the peptide nucleic acid is facing the 3' end of the nucleic acid sequence, and to hybridise in the parallel orientation when the C-terminal end of the peptide nucleic acid is facing the 5' end of the nucleic acid sequence. In most applications, hybridisation in the antiparallel orientation is preferred as the hybridisation in the parallel orientation takes place rather slowly and as the formed duplexes are not as stable as the duplexes having antiparallel strands. Triplex formation with a stoichiometry of two peptide nucleic acid strands and one nucleic acid strand may occur if the peptide nucleic acid has a high pyrimidine content. Such triplexes are very stable, and probes capable of forming triplexes may thus be suitable for certain applications.

Mainly because the peptide nucleic acid strand is uncharged, a peptide nucleic acid-nucleic acid-duplex will have a higher T_m than the corresponding nucleic acid-nucleic acid-duplex. Typically there will be an increase in T_m of about 1 °C per base pair at 100 mM NaCl depending on the sequence (Egholm et al. (1993), Nature, 365, 566-568).

In contrast to DNA-DNA-duplex formation, no salt is necessary to facilitate and stabilise the formation of a peptide nucleic acid-DNA or a peptide nucleic acid-RNA duplex. The T_m of the peptide nucleic acid-DNA-duplex changes only little with increasing ionic strength. Typically for a 15-mer, the T_m will drop only 5 °C when the salt concentration is raised from 10 mM NaCl to 1 M NaCl. At low ionic strength (e.g. 10 mM phosphate buffer with no salt added), hybridisation of a peptide nucleic acid to a target sequence is possible under conditions where no stable DNA-DNA-duplex formation occurs. Furthermore, target sites that normally are inaccessible can be made more readily accessible for hybridisation with peptide nucleic acid probes at low salt concentration as the secondary and tertiary structure of nucleic acids are destabilised under such conditions. Using peptide nucleic acid probes, a separate destabilising step or use of destabilising probes may not be necessary to perform.

rRNA is essential for proper function of the ribosomes and thus the synthesis of proteins. The genes encoding the rRNAs are in eubacteria located in an operon in which the small subunit RNA gene, the 16S rRNA gene, is located nearest the 5' end of the operon, the gene for the large subunit RNA, the 23S rRNA gene, is located distal to the 16S rRNA gene and the 5S rRNA gene is located nearest the 3' end of the operon. The three genes are separated by spacer regions in which tRNA genes may be found, however, there are none in *M. tuberculosis*. The primary transcript of the eubacterial rRNA operon is cleaved by RNase III. This cleavage results in separation of the 16S, the 23S and the 5S rRNA into precursor rRNA molecules (pre-rRNA molecules) which besides the rRNA species also contain leader and tail sequences. The primary RNase III cleavage is normally a rapid process, whereas the

subsequent maturation is substantially slower. Precursor rRNA is typically more abundant than even strongly expressed mRNA species. Thus, for certain applications, precursor rRNA may be an attractive diagnostic target. In order to specifically detect precursor rRNA, a target probe should be directed against sequences comprising at least part of the leader or tail
5 sequences. A target probe may further be directed against sequences of which both part of the leader/tail and mature rRNA sequences are constituents.

Usually, patients having contracted a mycobacterial infection are treated with medicaments until no mycobacteria can be found in the sputum. Except for culturing, the presently available
10 methods do not allow for clear distinguishing between living and dead mycobacteria. This means that a patient may often be treated with medicaments for a longer period of time than actually necessary. A way of determining the progress of treatment would be a very valuable tool in the fight of tuberculosis and other mycobacterial diseases.

15 As transcription and maturation of rRNA is a measure of viability, detection of precursor rRNA is a suitable and direct measure of viability of the bacteria. Furthermore, precursor rRNA may be used for identification of antibiotic drugs which reduce or inhibit rRNA transcription. One such example is rifampicin. A transcriptional inhibitor will in susceptible bacteria eliminate new synthesis of rRNA and thus the pool of precursor rRNA will be depleted. However, in resistant
20 cells, primary transcripts as well as precursor rRNAs will continue to be produced.

Although it is preferred to use peptide nucleic acid probes targeting specific sequences of rRNA, it will readily be understood that peptide nucleic acid probes complementary to rRNA targeting probes will be useful for the detection of the genes coding for said sequence specific
25 rRNA (rDNA), and peptide nucleic acid probes for the detecting rDNA is hence contemplated by the present invention. Although it is preferred to choose the sequence of the probe so as to enable the probe to hybridise to its target sequence in antiparallel orientation, it is to be understood that probes capable of hybridising in parallel orientation can be constructed from the same information. The present invention is intended to cover both types of probes.

30

In the broadest aspect, the present invention relates to peptide nucleic acid probes for detecting a target sequence of one or more mycobacteria optionally present in a test sample, said probe being capable of hybridising to a target sequence of mycobacterial rDNA, precursor rRNA or rRNA (claim 1).

35

The probes of the invention may suitably be directed to rDNA, precursor rRNA, or to 23S, 16S or 5S rRNA.

In accordance with claim 3, the target sequences, to which the peptide nucleic acid probe(s) are capable of hybridising to, are obtainable by

- 5 (a) comparing the nucleobase sequences of said mycobacterial rRNA or rDNA of one or more mycobacteria to be detected with the corresponding nucleobase sequence of organism(s), in particular other mycobacteria, from which said one or more mycobacteria are to be distinguished,
- (b) selecting a target sequence of said rRNA or rDNA which includes at least one nucleobase differing from the corresponding nucleobase of the organism(s), in particular other
- 10 mycobacteria, from which said one or more mycobacteria are to be distinguished, and
- (c) determining the capability of said probe to hybridise to the selected target sequence to form detectable hybrids.

Peptide nucleic acid probes are, in accordance with claim 4, obtainable by


15

- (a) comparing the nucleobase sequences of said mycobacterial rRNA or rDNA of one or more mycobacteria to be detected with the corresponding nucleobase sequence of organism(s), in particular other mycobacteria, in particular other mycobacteria, from which said one or more mycobacteria are to be distinguished,
- 20 (b) selecting a target sequence of said rRNA or rDNA which includes at least one nucleobase differing from the corresponding nucleobase of the organism(s), in particular other mycobacteria, from which said one or more mycobacteria are to be distinguished,
- (c) synthesising said probe, and
- (4) determining the capability of said probe to hybridise to the selected target sequence to
- 25 form detectable hybrids.

The probes are in particular suitable for detecting a target sequence of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC) or for detecting a target sequence of one or more mycobacteria other than mycobacteria of the Mycobacterium tuberculosis Complex (MOTT) optionally present in a sample, which probe comprises from 6

30 to 30 polymerised peptide nucleic acid moieties, said probe being capable of hybridising to a target sequence of mycobacterial rDNA, precursor rRNA or 23S, 16S or 5S rRNA forming detectable hybrids (claim 5). In accordance with claim 6, such probes may comprise peptide nucleic acid moieties of formula (I)

14



(I)

5

wherein each X and Y independently designate O or S,

each Z independently designates O, S, NR^1 , or $\text{C}(\text{R}^1)_2$, wherein each R^1 independently designate H, C_{1-6} alkyl, C_{1-6} alkenyl, C_{1-6} alkynyl,

10 each R^2 , R^3 and R^4 designate independently H, the side chain of a naturally occurring amino acid, the side chain of a non-naturally occurring amino acid, C_{1-4} alkyl, C_{1-4} alkenyl or C_{1-4} alkynyl, or a functional group, each Q independently designates a naturally occurring nucleobase, a non-naturally occurring nucleobase, an intercalator, a nucleobase-binding group, a label or H,

15 and with the proviso indicated in claim 6.

The probes may suitably be used for detecting a species specific mycobacterial target sequence, or target sequences of a group of mycobacteria like MTC, MOTT, MAC or MAIS.

20 The probes may further be designed so as to be capable of hybridising to one or more drug resistant mycobacteria, or, alternatively, to the wild-type corresponding thereto. In the design of the probes, sequences between different mycobacteria (one or more) may be taken into account as may sequences from other related or non-related organisms (one or more).

As mentioned above, drug-resistance is an increasing threat to the fight of mycobacterial infection. Monotherapy with macrolides such as clarithromycin and azithromycin often leads to clinically significant drug-resistance. Clarithromycin and azithromycin are important drugs in the treatment of infections by especially *M. avium*. Comparison between 23S rRNA sequences from isolates of *M. avium* and *M. intracellulare* with acquired resistance to clarithromycin and azithromycin and 23S rRNA sequences from non-resistant strains has revealed that the majority of resistant strains have single-point mutations in the 23S rRNA in positions corresponding to 2058 and 2059 in *E. coli* 23S rRNA. In the *M. avium* 23S rRNA sequence (GenBank accession number X74494), the corresponding bases are in position 2568 and 2569, respectively, as shown in Figure 6. Most susceptible strains have an A residue in one of these positions whereas the resistant strains have a C, G or T in position 2058 (*E. coli* numbering corresponding to 2568 in *M. avium* with GenBank accession number X74494), or G or C in position 2059 (*E. coli* numbering corresponding to 2569 in *M. avium* with GenBank accession number X74494).

Single-point mutations in rRNA apparently also account to some degree for streptomycin resistance. Streptomycin, the first successful antibiotic drug against tuberculosis, is an aminocyclitol glycoside that perturbs protein synthesis at the ribosomal level. The genetic basis for streptomycin resistance has not yet been completely solved. However, some
5 streptomycin resistant strains of *M. tuberculosis* have single-point mutations in 16S rRNA. These mutations are located in positions corresponding to bases 501, 522, 523, 526, 912 and 913 in *E. coli* 16S rRNA which correspond to bases with numbers 452, 473, 474, 477, 865 and 866, respectively, of *M. tuberculosis* 16S rRNA (GenBank accession number X52917) as shown in Figure 7. Most of these mutated nucleotides are involved in structural interactions
10 within the 530 loop of 16S rRNA which is one of the most conserved regions in the entire 16S rRNA gene.

Mutations in an 81 bp region of the gene (*rpoB*) encoding the beta subunit of RNA polymerase are the cause of 96% of the cases of rifampicin resistance in *M. tuberculosis* and *M. leprae*.
15 rRNA precursor molecules have terminal domains (tails) which are removed during late steps in precursor rRNA processing to yield the mature rRNA molecules. Transcriptional inhibitors such as rifampicin can deplete precursor rRNA in sensitive cells by inhibiting de novo precursor rRNA synthesis while allowing maturation to proceed. Thus, precursor rRNA is depleted in susceptible mycobacterium cells while it remains produced in resistant
20 mycobacterium cells when the cells are exposed to rifampicin during culturing.

Peptide nucleic acid probes for detecting a target sequence of one or more mycobacteria of the *Mycobacterium tuberculosis* Complex are defined in claims 7 to 10. Peptide nucleic acid probes for detecting a target sequence of one or more mycobacteria other than mycobacteria
25 of the *Mycobacterium tuberculosis* Complex are defined in claims 11 to 13. Peptide nucleic acid probes for detecting a target sequence of one or more drug resistant mycobacteria of the *Mycobacterium tuberculosis* complex or of one or more drug resistant mycobacteria other than mycobacteria of the *Mycobacterium tuberculosis* Complex are defined in claim 14.

30 In the present context and the claims, the term "naturally occurring nucleobases" includes the four main DNA bases (i.e. thymine (T), cytosine (C), adenine (A) and guanine (G)) as well as other naturally occurring nucleobases (e.g. uracil (U) and hypoxanthine).

The term "non-naturally occurring nucleobases" comprises i.a. modified naturally occurring
35 nucleobases. Such non-naturally occurring nucleobases may be modified by substitution by e.g. one or more C₁₋₈ alkyl, C₁₋₈ alkenyl or C₁₋₈ alkynyl groups or labels. Examples of non-naturally occurring nucleobases are purine, 2,6-diamino purine, 5-propynylcytosine (C propynyl), isocytosine (iso-C), 5-methyl-isocytosine (iso^{Me}C) (see e.g. Tetrahedron Letters Vol

36, No 12, 2033-2036 (1995) or Tetrahedron Letters Vol 36, No 21, 3601-3604 (1995)), 7-deazaadenine, 7-deazaguanine, N⁴-ethanocytosine, N⁶-ethano-2,6-diaminopurine, 5-(C₃₋₆)-alkenyluracil, 5-(C₃₋₆)-alkynylcytosine, 5-fluorouracil and pseudocytosine.

- 5 Examples of useful intercalators are e.g. acridin, anthraquinone, psoralen and pyrene.

Examples of useful nucleobase-binding groups are e.g. groups containing cyclic or heterocyclic rings. Non-limiting examples are 3-nitro pyrrole and 5-nitro indole.

- 10 It is to be understood that alkyl, alkenyl and alkynyl groups may be branched or non-branched, cyclic or non-cyclic, and may further be interrupted by one or more heteroatoms, or may be unsubstituted or substituted by or may contain one or more functional groups. Non-limiting examples of such functional groups are acetyl groups, acyl groups, amino groups, carbamido groups, carbamoyl groups, carbamyl groups, carbonyl groups, carboxy groups, cyano groups, dithio groups, formyl groups, guanidino groups, halogens, hydrazino groups, hydrazo groups, hydroxamino groups, hydroxy groups, keto groups, mercapto groups, nitro groups, phospho groups, phosphono groups, phospho ester groups, sulfo groups, thiocyanato groups, cyclic, aromatic and heterocyclic groups.
- 15
- 20 C₁₋₄ groups contain from 1 to 4 carbon atoms, C₁₋₆ groups contain from 1 to 6 carbon atoms, and C₁₋₁₅ groups contain from 1 to 15 carbon atoms, not including optional substituents, heteroatoms and/or functional groups. Non-limiting examples of such groups are -CH₃, -CF₃, -CH₂-, -CH₂CH₃, -CH₂CH₂-, -CH(CH₃)₂-, -OCH₃, -OCH₂-, -OCH₂CH₃, -OCH₂CH₂-, -OCH(CH₃)₂-, -OC(O)CH₃, -OC(O)CH₂-, -C(O)H, -C(O)-, -C(O)CH₃, -C(O)OH, -C(O)O-, -CH₂NH₂, -CH₂NH-,
- 25 -CH₂OCH₃, -CH₂OCH₂-, -CH₂OC(O)OH, -CH₂OC(O)O-, -CH₂C(O)CH₃, -CH₂C(O)CH₂-, -C(O)NH₂, -CH=CH₂, -CH=CH-, -CH=CHCH₂C(O)OH, -CH=CHCH₂C(O)O-, -C≡CH, -C≡C-, -CH₂C≡CH, -CH₂C≡C-, -CH₂C≡CCH₃, -OCH₂C≡CH, -OCH₂C≡C-, -OCH₂C≡CCH₃, -NHCH₂C(O)-, -NHCH₂CH₂C(O)-, -NH(CH₂CH₂O)₂CH₂C(O)-, and HO(O)CCH₂C(O)(NH-(CH₂CH₂O)₂CH₂C(O))₂-, phenyl, benzyl, naphthyl, oxazolyl, pyridinyl, thiadiazolyl, triazolyl,
- 30 and thienyl.

Within the present context, the expression "naturally occurring amino acid" is intended to comprise D- and L-forms of amino acids commonly found in nature, e.g. D- and L-forms of Ala (alanine), Arg (arginine), Asn (asparagine), Asp (aspartic acid), Cys (cysteine), Gln (glutamine),

35 Glu (glutamic acid), His (histidine), Ile (isoleucine), Leu (leucine), Lys (lysine), Met (methionine), Phe (phenylalanine), Pro (proline), Ser (serine), Thr (threonine), Trp (tryptophan), Tyr (tyrosine) and Val (valine).

In the present context, the expression "non-naturally occurring amino acid" is intended to comprise D- and L-forms of amino acids other than those commonly found in nature as well as modified naturally occurring amino acids. Examples of useful non-naturally occurring amino acids are D- and L-forms of β -Ala (β -alanine) Cha (cyclohexylalanine), Cit (citrulline), Hci (homocitrulline), HomoCys (homocystein), Hse (homoserine), Nle (norleucine), Nva (norvaline), Orn (ornithine), Sar (sarcosine) and Thi (thienylalanine).

In the present context, the term "sample" is intended to cover all types of samples suitable for the purpose of the invention. Examples of such samples are sputum, laryngeal swabs, gastric lavage, bronchial washings, biopsies, aspirates, expectorates, body fluids (spinal, pleural, pericardial, synovial, blood, pus, bone marrow), urine, tissue sections as well as food samples, soil, air and water samples. Analysis of samples originating from the before-mentioned samples (e.g. cultures and treated samples) are also within the scope of the invention.

In the present context, the term "hybrids" is intended to include complexes between a probe and a nucleic acid to be determined. Such hybrids may be made up of two or more strands.

The strength of the binding between the probe and the target nucleic acid sequence may be influenced by the ligand Q. When Q designates a nucleobase, Hoogsteen and/or Watson-Crick base pairing assist(s) in the formation of hybrids between a nucleic acid sequence to be detected and a probe. It is contemplated that one or more of the ligands may be a group which contribute little or none to the binding of the nucleic acid such as hydrogen. It is contemplated that suitable probes to be used comprise less than 25% by weight of peptide nucleic acid moieties, wherein Q designates such groups. One or more of the ligands Q may be groups that stabilise nucleobase stacking such as intercalators or nucleobase-binding groups.

In the above-indicated probes, one or more of the Q-groups may designate a label. Examples of suitable labels are given below. Moieties wherein Q denotes a label may preferably be located in one or both of the terminating moieties of the probe. Moieties wherein Q denotes a label may, however, also be located internally.

The peptide nucleic acid probes may comprise moieties, wherein all X groups are O (polyamides) or wherein all X groups are S (polythioamides). It is to be understood that the probes may also comprise mixed moieties (comprising both amide and thioamide moieties).


In another aspect, the present invention relates to peptide nucleic acid probes of formula (II), (III) and (IV) as well as mixtures of such probes defined in claim 15.

In a preferred embodiment, the peptide nucleic acid probes or mixtures thereof according to the invention are of formulas (I)-(IV) as defined in claim 16 with Z being NH, NCH₃ or O, each R², R³ and R⁴ independently being H or the side chain of a naturally occurring amino acid, the side chain of a non-naturally occurring amino acid, or C₁₋₄ alkyl, and each Q being a naturally occurring nucleobase or a non-naturally occurring nucleobase with the provisos defined in claims 6 to 14.

Peptide nucleic acid probes or mixtures of such probes according to the invention are preferably those of formula (I)-(IV) as defined in claim 17 with Z being NH or O, and R² being H or the side chain of Ala, Asp, Cys, Glu, His, HomoCys, Lys, Orn, Ser or Thr, and Q being a nucleobase selected from thymine, adenine, cytosine, guanine, uracil, iso-C, and 2,6-diaminopurine with the provisos defined in claims 6 to 14.

Peptide nucleic acid probes or mixtures thereof, which are of major interest for detecting mycobacteria of the MTC group or one or more mycobacteria other than mycobacteria of the MTC group, are probes of formula (V) according to claim 18, wherein R⁴ is H or the side chain of Ala, Asp, Cys, Glu, His, HomoCys, Lys, Orn, Ser or Thr, Q is as defined in claim 17 and with the provisos indicated in claims 6 to 14.

The peptide nucleic acid probe comprises polymerised moieties as defined above and in the claims. From the formula, it is to be understood that the probe may comprise polymerised moieties which structure may be mutually different or identical. In some cases, it may be advantageous that at least one moiety of the probe, preferably one (or both) of the moieties terminating the probe, are of a different structure. Such terminating moieties may suitably be a moiety of formula (VI)



(VI)

30

where Q is as defined above. Such moiety may suitably be connected to a peptide nucleic acid moiety through an amide bond.

The peptide nucleic acid probe according to the invention comprises from 6 to 30 polymerised moieties of formulas (I) to (V), and, in addition, optionally one or two terminating moieties of formula (VI) as defined above. The preferred length of the probe will depend on the sample material and whether labelled probes are used. It is contemplated that especially interesting probes comprise from 10 to 30 polymerised moieties of formulas (I) to (V), and, in addition,

optionally one or two terminating moieties of formula (VI) as defined above. Probes of the invention may suitably comprise from 12 to 25 polymerised moieties of formulas (I) to (V), more suitably from 14 to 22 polymerised moieties of formulas (I) to (V), most suitably from 15 to 20 polymerised moieties of formulas (I) to (V), and, in addition, optionally one or two
 5 terminating moieties of formula (VI).

As mentioned above, the polymerised moieties of the probes may be mutually different or identical. In some embodiments, the polymerised moieties of formulas (V) constitute at least 75% by weight (calculated by excluding labels and linkers), preferably at least 80% by weight
 10 and most preferably at least 90% by weight of the probe.

The ends on the moieties terminating the probe may be substituted by suitable substituents which in the following will be named "linkers". A terminating end may suitably be substituted by from 1 to 5 linkers, more suitably from 1 to 3 linkers. Such linkers may suitably be selected
 15 among C₁₋₁₅ alkyl, C₁₋₁₅ alkenyl and C₁₋₁₅ alkynyl groups as defined above. The linkers may be substituted or unsubstituted, branched or non-branched, or be interrupted by heteroatoms, or be substituted or contain functional groups as described above. This may depend on the chemical nature of the terminating moiety (i.e. whether the moiety is terminated by a carbon, oxygen or nitrogen atom). A terminating end or a linker on a terminating end may further be
 20 substituted by one or more labels, which labels may be incorporated end to end, i.e. so as to form a non-branched labelled end, or may be incorporated so as to form a branched labelled end ("zipper"). The linkers may be attached directly to a terminating end, may be attached to a label or between labels on a terminating end, or be attached to a terminating end before a label is attached to a terminating end. It should be understood that two terminating ends may
 25 carry different or identical substituents, linkers and/or labels. It should further be understood that the term "a label" is intended to comprise one or more labels as the term "linkers" is to comprise one or more linkers. For certain applications, it may be advantageous that one or more linkers are incorporated between the peptide nucleic acid moieties. Such applications may in particular be those based on triplex formation.

30

Examples of suitable linkers are -NH(CH₂CH₂O)_nCH₂C(O)-, -NH(CHOH)_nC(O)-, -(O)C(CH₂OCH₂)_nC(O)- and -NH(CH₂)_nC(O)-, NH₂(CH₂CH₂O)_nCH₂C(O)-, NH₂(CHOH)_nC(O)-, HO(O)C(CH₂OCH₂)_nC(O)-, NH₂(CH₂)_nC(O)-, -NH(CH₂CH₂O)_nCH₂C(O)OH, -NH(CHOH)_nC(O)OH, -(O)C(CH₂OCH₂)_nC(O)OH and -NH(CH₂)_nC(O)OH, wherein n is 0 or an
 35 integer from 1 to 8, preferably from 1 to 3. Examples of very interesting linkers are -NHCH₂C(O)-, -NHCH₂CH₂C(O)-, -NH(CH₂CH₂O)₂CH₂C(O)-, and HO(O)CCH₂CH₂C(O)(NH(CH₂CH₂O)₂CH₂C(O))₂-.

In the present context, the term "label" refers to a substituent which is useful for detection or visualisation. Suitable labels comprise fluorophores, biotin, dinitro benzoic acid, digoxigenin, radioisotope labels, peptide or enzyme labels, chemiluminescence labels, fluorescent particles, hapten, antigen or antibody labels.

5

The expression "peptide label" is intended to mean a label comprising from 1 to 20 naturally occurring or non-naturally occurring amino acids, preferably from 1 to 10 naturally occurring or non-naturally occurring amino acids, more preferably from 1 to 8 naturally occurring or non-naturally occurring amino acids, most preferably from 1 to 4 naturally occurring or non-naturally occurring amino acids, linked together end to end in a non-branched or branched ("zipper") fashion. Such peptide label may be composed of amino acids which are mutually identical or different. In a preferred embodiment, such a non-branched or branched end comprises one or more, preferably from 1 to 8 labels, more preferably from 1 to 4, most preferably 1 or 2, further labels other than a peptide label. Such further labels may suitably terminate a non-branched end or a branched end. One or more linkers may suitably be attached to the terminating end before a peptide label and/or a further label is attached. Such linker units may also be attached between a peptide label and a further label. Furthermore, such peptide labels may be incorporated between the peptide nucleic acid moieties.

20 The probe as such may also comprise one or more labels such as from 1 to 8, preferably from 1 to 4, most preferably 1 or 2, labels and/or one or more linker units, which may be attached internally, i.e. to the backbone of the probe. The linker units and labels may mutually be attached as described above.

25 Examples of particular interesting labels are biotin, fluorescein labels, e.g. 5-(and 6)-carboxy-fluorescein, 5- or 6-carboxyfluorescein, 6-(fluorescein)-5-(and 6)-carboxamido hexanoic acid and fluorescein isothiocyanate, peptide labels consisting of from 1 to 20 naturally occurring amino acids or non-naturally occurring amino acids, enzyme labels such as peroxidases like horse radish peroxidase (HRP), alkaline phosphatase, and soya bean peroxidase, dinitro benzoic acid, rhodamine, tetramethylrhodamine, cyanine dyes such as Cy2, Cy3 and Cy5, coumarin, R-phycoerythrin (RPE), allophycoerythrin, Texas Red, Princeton Red, and Oregon Green as well as conjugates of R-phycoerythrin and, e.g. Cy5 or Texas Red.

35 Examples of preferred labels are biotin, fluorescent labels, peptide labels, enzyme labels and dinitro benzoic acid. Peptide labels may preferably be composed of from 1 to 10, more preferably of from 1 to 8, most preferably of from 1 to 4, naturally occurring or non-naturally occurring amino acids. It may be particularly advantageous to incorporate one or more other labels as well as a peptide label such as from 1 to 8 or from 1 to 4 other labels, preferably 1 or

2 other labels.

Suitable peptide labels may preferably be composed of cysteine, glycine, lysine or ornithine.

- 5 In a further embodiment, the Q substituent as defined above may be labelled. Suitable labels are as defined above. Between Q and such a label, a linker as defined above may be incorporated. It is preferred that such labelled ligands Q are selected from thymine and uracil labelled in the 5-position and 7-deazaguanine and 7-deazaadenine labelled in the 7-position.
- 10 A mixture of peptide nucleic acid probes is also part of the present invention. Such mixture may comprise more than one probe capable of hybridising to 23S rRNA, and/or more than one probe capable of hybridising to 16S rRNA, and/or or more than one probe capable of hybridising to 5S rRNA. A mixture of probes may further comprise probe(s) directed to precursor rRNA and/or rDNA. The mixture may also comprise peptide nucleic acids for
- 15 detecting more than one mycobacteria in the same assay.

In a preferred embodiment, the nucleobase sequence of the peptide nucleic acid probe is selected so as to be substantially complementary to the nucleobase sequence of the target sequence in question. In an especially preferred embodiment, the nucleobase sequence of the

20 peptide nucleic acid probe is selected so as to be complementary to the nucleobase sequence of the target sequence in question. By "complementary" is meant that the nucleobases are selected so as to enable perfect match between the nucleobases of the probe and the nucleobases of the target, i.e. A to T or G to C. By substantially complementary is meant that the peptide nucleic acid probe is capable of hybridising to the target sequence, however, the

25 probe does not necessarily have to be perfectly complementary to the target. For example, probes comprising one or more bases not complementary to the target sequence and non-target sequences, especially base(s) located at the end of the probe, where the effect on the stability of probe-target nucleic acid hybrids is low. Another example is probes comprising other naturally occurring bases. Thus provided that the probe is capable of hybridising to the

30 target sequence, the nucleobase difference(s) between target sequences and non-target sequences ensures that the stability of probe-non-target nucleic acid hybrids are lower than the stability of probe-target nucleic acid hybrids and therefore make such substantially complementary probes applicable for detection of mycobacteria.

- 35 The probes may be synthesised according to the procedures described in "PNA Information Package" obtained from Millipore Corporation (Bedford, MA, USA), or may be synthesised on an Expedite Nucleic Acid Synthesis System (PerSeptive BioSystems, USA).

If using the Fmoc strategy for elongation of the probe with linkers or amino acids, it is possible to retain side chain amino groups protected with acid sensitive protection groups such as the Boc or Mtt group. This method allows introduction of a linker containing several Boc protected amino groups which can all be cleaved and labelled in the same synthesis cycle.

5

One way of labelling a probe is to use a fluorescent label, such as 5-(and 6)-carboxyfluorescein, 5- or 6-carboxyfluorescein, or 6-(fluorescein)-5-(and 6)-carboxamido hexanoic acid. The acid group is activated with HATU (O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate) or HBTU (2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate) and reacted with the N-terminal amino group of the peptide nucleic acid. The same technique can be applied to other labelling groups containing an acid function. Alternatively, the succinimidyl ester of the above-mentioned labels may suitably be used or fluorescein isothiocyanate may be used directly.

10

After synthesis, probes can be cleaved from the resin using standard procedures as described by Millipore Corporation or PerSeptive BioSystems. The probes are subsequently purified and analysed using reversed-phase HPLC techniques at 50°C and were characterised by matrix-assisted laser desorption/ionisation time of flight mass spectrometry (MALDI-TOFMS), plasma desorption mass spectrometry (PDMS), electron spray mass spectrometry (ESMS), or fast atom bombardment (FAB-MS).

20

Generally, probes such as probes comprising polymerised moieties of formula (IV) and (V) may also be prepared as described in, e.g., *Angewandte Chemie, International Edition in English* 35, 1939-1942 (1996) and *Bioorganic & Medical Chemistry Letters*, Vol 4, No 8, 1077-1080 (1994). Chemical properties of some probes are described in, e.g., *Nature*, 365, 566-568 (1993).

25

The method as claimed can be used for the detection of a target sequence of one or more mycobacteria optionally present in a sample. The method and the probes provide a valuable tool for analysing samples for the presence of such target sequences, hence providing information for establishing a diagnosis.

30

In the assay method according to the invention, the sample to be analysed for the presence of mycobacteria is brought into contact with one or more probes or a mixture of such probes according to the invention under conditions by which hybridisation between the probe(s) and any sample rRNA or rDNA originating from mycobacteria can occur, and the formed hybrids, if any, are observed or measured, and the observation or measurement is related to the presence of a target sequence of one or more mycobacteria. The observation or

35

measurement may be accomplished visually or by means of instrumentation.

Prior to contact with probe(s) according to the invention, the samples may undergo various types of sample processing which include purification, decontamination and/or concentration.

- 5 The sample may suitably be decontaminated by treatment with sodium hypochlorite and subsequently centrifuged for concentration of the mycobacteria. Samples e.g. sputum samples may be treated with a mucolytic agent such as N-Acetyl-L-cystein which reduces the viscosity of the sample as well as be treated with sodium hydroxide which decontaminates the sample, and subsequently centrifuged. Other well-known decontamination and concentration
- 10 procedures include the Zephiran-trisodium phosphate method, Petroff's sodium hydroxide method, the oxalic acid method as well as the cetylpyridinium chloride-sodium chloride method. Samples may also be purified and concentrated by applying sample preparation methods such as filtration, immunocapture, two-phase separation either alone or in combination. The sample preparation methods may also be used together with the
- 15 centrifugation and decontamination methods mentioned above.

- Samples may, either directly or after having undergone one or more processing steps, be analysed in primarily two major types of assays, in situ-based assays and in vitro-based assays. In this context, in situ-based assays are to be understood as assays, in which the
- 20 target nucleic acids remain within the bacterial cell during the hybridisation process. Examples are in situ hybridisation (ISH) assays on smears and biopsies as well as hybridisation to whole cells which may be in suspension and which subsequently may be analysed by e.g. flow cytometry optionally after capture of the bacteria onto particles (with same or different type and size), or by image analysis after spreading of the bacteria onto a solid medium, filter
- 25 membrane or another substantially planar surface.

- In vitro-based assays are to be understood as assays, in which the target nucleic acids are released from the bacterial cell before hybridisation. Examples of such assays are microtiter plate-based assays. Many other assay types, in which the released target nucleic acids by
- 30 some means are captured onto a solid phase and subsequently analysed via a detector probe, are feasible and within the scope of the present invention. Even further, in vitro-based assays include assays, in which the target nucleic acids are not captured onto a solid phase, but in which the hybridisation and signal generation take place entirely in solution.

- 35 Samples for in situ-based assays may suitably be applied and optionally be immobilised to a support. Techniques for applying of a sample onto a solid support depend on the type of sample in question and include smearing and cytocentrifugation for liquid or liquified samples and sectioning of tissues for biopsy materials. The solid support may take a wide variety of

forms well-known in the art, such as a microscope slide, a filter membrane, a polymer membrane or a plate of various materials.

- In the case of in vitro-based assays, the target nucleic acid may be released from the mycobacterial cells in various ways. Most methods for releasing the nucleic acids cause bursting of the cell wall (lysis) followed by extraction of the nucleic acids into a buffered solution. As mycobacteria have complex cell walls containing covalently associated peptidoglycans, arabinogalactans and in particular mycolic acids, they cannot easily be disrupted by standard methods used for the rapid lysis of other bacteria. Possible methods which are known to give successful lysis of the mycobacterial cell wall include methods which involve treatment with organic solvents, treatment with strong chaotropic reagents such as high concentrations of guanidine thiocyanate, enzyme treatment, bead beating, heat treatment, sonication and/or application of a French press.
- Samples to be analysed by in situ assays may be fixed prior to hybridisation. The person skilled in the art will readily recognise that the appropriate procedure will depend on the type of sample to be examined. Fixation and/or immobilisation should preferably preserve the morphological integrity of the cellular matrix and of the nucleic acids. Examples of methods for fixation are flame fixation, heat fixation, chemical fixation and freezing. Flame fixation may be accomplished by passing the slide through the blue cone of a Bunsen burner 3 or 4 times; heat fixation may be accomplished by heating the sample to 80°C for 2 hours; chemical fixation may be accomplished by immersion of the sample in a fixative (e.g. formamide, methanol or ethanol). Freezing is particularly relevant for biopsies and tissue sections and is usually carried out in liquid nitrogen.
- In one in situ hybridisation assay embodiment, the sample to be analysed is smeared onto a substantially planar solid support which may be a microscope slide, a filter membrane, a polymer membrane or another type of solid support with a planar surface. The preferred solid support is a microscope slide. After the smear has been prepared, it may optionally undergo further pre-treatment like treatment with bactericidal agents or additional fixation by immersion in e.g. ethanol. The sample may also be pre-treated with enzyme(s) which as primary function permeabilise the cells and/or reduce the viscosity of the sample. It may further be advantageous to perform a pre-hybridisation step in order to block sites which might otherwise give rise to non-specific binding. For this purpose, blocking agents like skim milk, and non-target probes may suitably be used. The components of the pre-hybridisation mixture should be selected so as to obtain an effective saturation of sites in the sample that might otherwise bind the probe non-specifically. The pre-hybridisation buffer may suitably comprise an appropriate buffer, blocking agent(s), and detergents.

During the in situ hybridisation, one or more probes according to the present invention are brought into contact with any target rRNA or rDNA inside the cells in a hybridisation solution under suitable stringency conditions. The concentration of the applied probe may vary
5 depending on the chemical nature of the probe and the conditions under which hybridisation is carried out. Typically, a probe concentration between 1 nM and 1 μ M is suitable. The hybridisation solution may comprise a denaturing agent which allows hybridisation to take place at a lower temperature than would be the case without the agent. The denaturing agent should be present in an amount effective to increase the ratio between specific binding and
10 non-specific binding. The effective amount of denaturing agent depends on the type used and on the probe or combination of probes. Examples of denaturing agents are formamide, ethylene glycol and glycerol, and these may preferably be used in a concentration above 10% and less than 70%. The preferred denaturing agent is formamide which is used more preferably in concentrations from 20% to 60%, most preferably from 30% to 50%. It should be
15 noted that in several instances it may not be necessary or advantageous to include a denaturing agent.

Prior to hybridisation or during hybridisation, a mixture of random probes (probes with random, non-selected sequences of optionally different length) may be added in excess to reduce non-
20 specific binding. Also, one or more non-sense probes (probes with a defined nucleobase sequence and length differing from the nucleobase sequence of the target sequence) may be added in excess in order to reduce non-specific binding. Also, non-specific binding of detectable probes to one or more non-target nucleic acid sequences can be suppressed by addition of one or more unlabelled or independently detectable probes, which probes have a
25 sequence that is complementary to the non-target sequence(s). It is particularly advantageous to add such blocking probes when the non-target sequence differs from the target sequence by only one mismatch.

It may be advantageous to include inert polymers which are believed to increase the effective
30 concentration of the probe(s) in the hybridisation solution. One such macromolecule is dextran sulphate which may be used in concentrations of from 2.5% to 15%. The preferred concentration range is from 8% to 12% in the case of dextran sulphate. Other useful macromolecules are polyvinylpyrrolidone and ficoll, which typically are used at lower concentrations, e.g. 0.2%. It may further be advantageous to add one or more detergents
35 which may reduce the degree of non-specific binding of the peptide nucleic acid probes. Examples of useful detergents are sodium dodecyl sulphate, Tween 20[®] or Triton X-100[®]. Detergents are usually used in concentrations between 0.05% and 1.0%, preferably between 0.05% and 0.25%. The hybridisation solution may furthermore contain salt. Although it is not

necessary to include salt in order to obtain proper hybridisation, it may be advantageous to include salt in concentrations from 2 to 500 mM, or suitably from 5 to 100 mM.

During hybridisation, other important parameters are hybridisation temperature, concentration
5 of the probe and hybridisation time. The person skilled in the art will readily recognise that optimal conditions must be determined for each of the above-mentioned parameters according to the specific situation, e.g. choice of probe(s) and type and concentration of the components of the hybridisation buffer, in particular the concentration of denaturing agent. Presence of volume excluders may also have an effect.

10

Following hybridisation, the sample is washed to remove any unbound and any non-specifically bound probe, and consequently, appropriate stringency conditions should be used. By stringency is meant the degree to which the reaction conditions favour the dissociation of the formed hybrids. The stringency may be increased typically by increasing the washing
15 temperature and/or washing time. Typically, washing times from 5 to 40 minutes may be sufficient. Two or more washing periods of shorter time may also give good results. A range of buffers may be used, including biological buffers, phosphate buffers and standard citrate buffers. The demand for low salt concentration in the buffers is not as pertinent as for DNA probe assays due to the difference response to salt concentration. In some cases, it is
20 advantageous to increase the pH of the washing buffer as it may give an increased signal-to-noise ratio (see WO 97/18325). This is conceivably due to a significant reduction of the non-specific binding. Thus, it may be advantageous to use a washing solution with a pH value from 8 to 10.5, preferably from 9 to 10.

25 Visualisation of bound probe(s) must be carried out with due regard to the type of label chosen. There are a wide range of useful probe labels, in particular various fluorescent labels such as fluorescein, rhodamine and derivatives thereof. Furthermore, labels like enzymes (e.g. peroxidases and phosphatases) and haptens (e.g. biotin, digoxigenin, dinitro benzoic acid) may suitably be applied. In the case of fluorescent labels, the hybrids formed may be
30 visualised using a microscope with a magnification of at least $\times 250$, preferably $\times 1000$. The visualisation may further be carried out using a CCD (charge coupled device) camera optionally controlled by a computer. When haptens are used as labels, the hybrids may be detected using an antibody conjugated with an enzyme. In these cases, the detection step may be based on colorimetry, fluorescence or luminescence.

35

The probes may alternatively be labelled with fluorescent particles having the fluorescent label embedded in the particles (e.g. Estapor K coloured microspheres), located on the surface of the particles and/or coupled to the surfaces of the particles. As the particles have to come into

contact with the target nucleic acids within the cells, the use of fluorescent particles may necessitate pretreatment of the bacteria. Relatively small particles e.g. about 20 nm may be suitable.

- 5 In another in situ hybridisation embodiment, frozen tissue or biopsy samples are cut into thin sections and transferred to a substantially planar surface, preferably microscope slides. Prior to hybridisation, the tissue or biopsy may be treated with a fixative, preferably a precipitating fixative such as acetone, or the sample is incubated in a solution of buffered formaldehyde. Alternatively, the biopsy or tissue section can be transferred to a fixative such as buffered
- 10 formaldehyde for 12 to 24 hours and following fixation, the tissue may be embedded in paraffin forming a block from which thin sections can be cut. Prior to hybridisation, the tissue section is dewaxed and rehydrated using standard procedures. Permeabilisation (e.g. treatment with proteases, diluted acids, detergents, alcohol and/or heat) may in some cases be advantageous. The selected method for permeabilisation depends on several factors, for
- 15 instance on the fixative used, the extent of fixation, the type and size of sample, and on the applied probe. For these types of samples, sample processing, prehybridisation, hybridisation, washing and visualisation may be carried out using same or adjusted conditions as described above.
- 20 In a further embodiment of the in situ assays, the bacterial cells are kept in suspension during fixation, prehybridisation, hybridisation and washing are carried out under the same or similar conditions as described above. The preferred type of label for this embodiment is fluorescent labels. This allows detection of hybridised cells by flow cytometry, recording the intensity of fluorescence per cell. Bacterial cells in suspension may further be coupled to particles,
- 25 preferably with a size of from 20 nm to 10 μ m. The particles may be made of materials well-known in the art like latex, dextran, cellulose and/or agarose, and may optionally be paramagnetic or contain a fluorescent label. Normally, bacterial cells are coupled to particles using antibodies against the target bacteria, but other means like molecular imprinting may also be used. Coupling of the bacterial cells to particles may be advantageous in sample
- 30 handling and/or during detection.

- In the embodiments of in situ hybridisation described above, the probes according to the invention are used for detecting a target sequence of one or more mycobacteria. In a preferred embodiment, the probes are suitable for detecting a target sequence of
- 35 mycobacteria of the Mycobacterium tuberculosis Complex (MTC), mycobacteria other than the Mycobacterium tuberculosis Complex (MOTT), or mycobacteria of the Mycobacterium avium Complex (MAC). The probes are further suitable for detecting simultaneously different target sequences originating from the same mycobacteria.

Samples to be analysed using in vitro-based assays need to undergo a treatment by which the nucleic acids are released from the bacterial cells. Nucleic acids may be released using organic solvents, strong chaotropic reagents such as high concentrations of guanidine thiocyanate, enzymes, bead beating, heating, sonication and/or application of a French press. The obtained nucleic acids may undergo additional purification prior to hybridisation.

In one in vitro hybridisation embodiment, the sample comprising the target nucleic acid is added to a container comprising immobilised capture probe(s) and one or more probe(s) labelled to function as detector probe(s). The hybridisation should be performed under suitable stringency conditions. The hybridisation solution may further comprise a denaturing agent, blocking probes, inert polymers, detergents and salt as described for the in situ-type assays. Likewise, the hybridisation temperature, probe concentration and hybridisation time are important parameters that need to be controlled according to the specific conditions of the assay, e.g. choice of peptide nucleic acid probe(s) and concentration of some of the ingredients of the hybridisation buffer. If hybridisation of the target nucleic acid to the capture probe(s) and detector probe(s), respectively, is performed in two separate steps, different parameters, in particular different stringency conditions, may be used in these steps. The concentration of the capture probe may be higher for in situ assays as hybridisation may be controlled better and washing can be performed more efficiently.

The capture probes may be immobilised onto a solid support by any means, e.g. by a coupling reaction between a carboxylic acid on a linker and an amino derivatised support. The capture probe may further be coupled onto the solid support by photochemical activation of photoreactive groups which have been attached absorptively to the solid support prior to photochemical activation. Such photoreactive groups are described in the US 5 316 784 A. The capture probes may further be coupled to a hapten which allows an affinity based immobilisation to the solid support. One such example is coupling of a biotin to the probe(s) and immobilisation via binding to a streptavidin-coated surface.

The solid support may take a wide variety of forms well-known in the art, such as a microtiter plate having one or more wells, a filter membrane, a polymer membrane, a tube, a dip stick, a strip and particles. Filter membranes may be made of cellulose, celluloseacetate, polyvinylidene fluoride or any other materials well-known in the art. The polymer membranes may be of polystyrene, nylon, polypropylene or any other materials well known in the art. Particles may be paramagnetic beads, beads made of polystyrene, polypropylene, polyethylene, dextran, nylon, amyloses, celluloses, polyacrylamides and agarose. When the solid support has the form of a filter, a membrane, a strip or beads, it (they) may be

incorporated into a single-use device.

The selection of the label of the detector probe(s) depend on the specific assay format and possible instrumentation. When biotin labelled probes are used, the hybrids may be detected using streptavidin or an antibody against the biotin label which antibody or streptavidin may be conjugated with an enzyme and the actual detection depend on the choice of the specific enzyme, preferably a phosphatase or a peroxidase, and the substrate for the selected enzyme. The signal may in some cases be enhanced using commercially available amplification systems such as the catalysed signal amplification system for biotinylates probes (CSA by DAKO). Various polymer-based enhancement systems may also be used. An example is a dextran polymer to which both a hapten specific antibody and an enzyme is coupled. The detector probe(s) may further be labelled with other haptens, e.g. digoxigenin, dinitro benzoic acid and fluorescein, in which case the hybrids may be detected using an antibody against the hapten which antibody may be conjugated with an enzyme. It is even possible to apply detector probe(s) which have enzymes coupled directly onto the probes. There are a wide range of possibilities for selection of enzyme substrates allowing for colourimetric (substrates e.g. p-nitro-phenyl phosphate or tetra-methyl-benzidine), fluorogenic (substrates e.g. 4-methylumbilliferylphosphate) or chemiluminescent (substrates e.g. 1,2-dioxetanes) detection.

20

The detector probes may further be labelled with various fluorescent labels, preferably fluorescein or rhodamine, in which case the hybrids may be detected by measuring the fluorescence.

25 The detector probe(s) will typically be different from the capture probe(s), thus ensuring dual species specificity. The dual specificity will most often allow at least one of the probes to be shorter, e.g. a 10 mer probe.

Furthermore, the capture of purine rich sequences may be improved by utilising bis-peptide nucleic acids as capture probes. Such bis-peptide nucleic acids are described in WO 96/02558. The bis-peptide nucleic acids comprise a first peptide nucleic acid strand capable of hybridising in parallel fashion to the target nucleic acid, and a second peptide nucleic acid strand capable of hybridising in antiparallel fashion to the purine rich sequence of the nucleic acid to be captured. The two peptide nucleic acid strands are connected by a linker and are in this way capable of forming a triplex structure with said purine rich sequence nucleic acid. The number of polymerised moieties of each linker-separated peptide nucleic acid may be as previously defined for non-bis-peptide nucleic acids. However, due to the high stability of the triplexes formed, bis-peptide nucleic acids with short first and second strands can be used

30

35

making the design of a pyrimidine rich probe easier.

Instead of using a detector probe, capture probe: nucleic acid complexes may be detected using a detection system based on an antibody reacting specifically with complexes formed
5 between peptide nucleic acids and nucleic acids (such as described in WO 95/17430), in which detection system the primary antibody may comprise a label, or which detection system comprises a labelled secondary antibody, which specifically binds to the primary antibody. The specific detection again depends on the selected substrate which may be of any type of those mentioned above.

10

Depending on the type of specific assay format, label and detection principle various types of instrumentation may be used including conventional microplate readers, luminometers and flow cytometers. Adaptation of adequate instrumentation may allow for automatisisation of the assay.

15

In an example of this embodiment, a capture probe of the present invention is coupled to a microtiter plate by a photochemical reaction between anthraquinon-labelled capture probe and polystyrene of the microwell. Target rRNA is added to the microwells and incubated under stringent conditions. Unbound rRNA is removed by washing and the microwell are incubated
20 with a hapten-labelled detector probe under stringent conditions. The visualisation is carried out using an enzyme-labelled antibody against the hapten, which after removal of unbound antibody is detected using a chemiluminescence substrate.

In another example of this embodiment capture probes are coupled to latex particles, and
25 hybridisation is carried out under suitable conditions in the presence of e.g. fluorescein labelled detector probe(s). After hybridisation and optionally washing, the hybrids are detected by flow cytometry. A range of different beads (e.g. by size or colours) may carry different capture probes for different targets, thus allowing a multiple detection system.

30 In a further embodiment of the in vitro assays format, the capture probe, the target nucleic acid and the detector probe may hybridise in solution, and subsequently the capture probe is attached to a solid phase. The solid phase, the hybridisation conditions and means of detection may be selected according to the specific method as described above.

35 In a further embodiment of in vitro assays, the target nucleic acid may be immobilised onto filter or polymer membranes or other types of solid phases well-known in the art. The hybridisation conditions and means of detection may be selected according to the specific set-up as described above.

In a further embodiment of the in vitro assay, an array of up to 100 or even more different probes directed against different target sequences may be immobilised onto a solid surface and hybridisation of the target sequences to all the probes is carried out simultaneously. The solid phase, the hybridisation conditions and means of detection may be as described above. This allow for simultaneous detection or identification of a range of parameters, i.e. species identification and resistance patterns.

The present probes further provide a method of diagnosing infection by mycobacteria and a method for determining the stage of the infection and the appropriate treatment by which methods one or more optionally labelled probes according to the invention are brought into contact with a patient sample and the type of treatment and/or the effect of a treatment is (are) evaluated.

Kits comprising at least one peptide nucleic acid probe as defined herein are also part of the present invention. Such kit may further comprise a detection system with at least one detecting reagent and/or a solid phase capture system.

DESCRIPTION OF SPECIFIC EMBODIMENTS

Examples of suitable Qs of adjacent moieties are given below. Peptide nucleic acid probes comprising such Qs will be suitable for detecting mycobacteria, in particular mycobacteria of the MTC group or mycobacteria other than mycobacteria of the MTC group. The probes are written from left to right corresponding to from the N-terminal end towards the C-terminal end. Suitable Q subsequences for detecting 23S and 16S rRNA as well as 5S rRNA of the MTC group are given below. Suitable Q subsequences for detecting 23S and 16S rRNA of mycobacteria other than mycobacteria of the MTC group are further given below. The Q subsequences include at least one nucleobase complementary to a nucleobase selected from the positions given in parenthesis. The Q subsequences are given as non-limiting examples of construction of suitable probe nucleobase sequences. It is to be understood that the probes may comprise fewer or more peptide nucleic acid moieties than indicated.

MTC group (23S)

AGA TGC GGG TAG CAC (selected from positions 149-158 in Figure 1A),	(Seq ID no 1)
TGT TTT CTC CTC CTA (selected from positions 220-221 in Figure 1A),	(Seq ID no 2)
ACT GCC TCT CAG CCG (selected from positions 328-361 in	
Figure 1A and Figure 1B),	(Seq ID no 3)
TGA TAC TAG GCA GGT (selected from positions 453-455 in Figure 1B),	(Seq ID no 4)
CGG ATT CAC AGC GGA (selected from positions 490-501 in Figure 1B),	(Seq ID no 5)

	TCA CCA CCC TCC TCC (selected from positions 637-660 in Figure 1C),	(Seq ID no 6)
	CCA CCC TCC TCC (selected from positions 637-660 in Figure 1C)	(modified Seq ID no 6)
	TTA ACC TTG CGA CAT (selected from positions 706-712 in Figure 1D),	(Seq ID no 7)
	ACT ATT CAC ACG CGC (selected from positions 762-789 in Figure 1D),	(Seq ID no 8)
5	CTC CGC GGT GAA CCA (selected from position 989 in Figure 1D),	(Seq ID no 9)
	GCT TTA CAC CAC GGC (selected from positions 1068-1072 in Figure 1E),	(Seq ID no 10)
	ACG CTT GGG GGC CTT (selected from position 1148 in Figure 1E),	(Seq ID no 11)
	CCA CAC CCA CCA CAA (selected from positions 1311-1329 in Figure 1E),	(Seq ID no 12)
	CCG GTG GCT TCG CTG (selected from positions 1361-1364 in Figure 1F),	(Seq ID no 13)
10	ACT TGC CTT GTC GCT (selected from position 1418 in Figure 1F),	(Seq ID no 14)
	GAT TCG TCA CGG GCG (selected from positions 1563-1570 in Figure 1F),	(Seq ID no 15)
	AAC TCC ACA CCC CCG (selected from positions 1627-1638 in Figure 1G),	(Seq ID no 16)
	ACT CCA CAC CCC CGA (selected from positions 1627-1638 in Figure 1G),	(Seq ID no 17)
	ACC CCT TCG CTT GAC (selected from positions 1675-1677 in Figure 1G),	(Seq ID no 18)
15	CTT GCC CCA GTG TTA (selected from position 1718 in Figure 1G),	(Seq ID no 19)
	CTC TCC CTA CCG GCT (selected from positions 1734-1740 in Figure 1H),	(Seq ID no 20)
	GAT ATT CCG GTC CCC (selected from positions 1967-1976 in Figure 1H),	(Seq ID no 21)
	ACT CCG CCC CAA CTG (selected from positions 2403-2420 in Figure 1H),	(Seq ID no 22)
	CTG TCC CTA AAC CCG (selected from positions 2457-2488 in Figure 1I),	(Seq ID no 23)
20	TTC GAG GTT AGA TGC (selected from positions 2457-2488 in Figure 1I),	(Seq ID no 24)
	GTC CCT AAA CCC GAT (selected from positions 2457-2488 in Figure 1I),	(Seq ID no 25)
	GGT GCA CCA GAG GTT (selected from positions 2952-2956 in Figure 1I),	(Seq ID no 26)
	CTG GCG GGA CAA CTG (selected from positions 2966-2969 in Figure 1J),	(Seq ID no 27)
	TTA TCC TGA CCG AAC (selected from positions 3000-3003 in Figure 1J),	(Seq ID no 28)
25	GAC CTA TTG AAC CCG (selected from positions 3097-3106 in Figure 1J),	(Seq ID no 29)

MTC group (16S)

	GAA GAG ACC TTT CCG (selected from positions 76-79 in Figure 2A),	(Seq ID no 30)
	CAC TCG AGT ATC TCC (selected from positions 98-101 in Figure 2A),	(Seq ID no 31)
30	ATC ACC CAC GTG TTA (selected from positions 136-136 in Figure 2A),	(Seq ID no 32)
	GCA TCC CGT GGT CCT (selected from positions 194-201 in Figure 2B),	(Seq ID no 33)
	CAC AAG ACA TGC ATC (selected from positions 194-201 in Figure 2B),	(Seq ID no 34)
	TAA AGC GCT TTC CAC (selected from positions 222-229 in Figure 2B),	(Seq ID no 35)
	GCT CAT CCC ACA CCG (selected from position 242 in Figure 2B),	(Seq ID no 36)
35	CCG AGA GAA CCC GGA (selected from position 474 in Figure 2C),	(Seq ID no 37)
	AGT CCC CAC CAT TAC (selected from positions 1136-1145 in Figure 2C),	(Seq ID no 38)
	AAC CTC GCG GCA TCG (selected from positions 1271-1272 in Figure 2C),	(Seq ID no 39)
	GGC TTT TAA GGA TTC (selected from positions 1287-1292 in Figure 2D),	(Seq ID no 40)
	GAC CCC GAT CCG AAC (selected from position 1313 in Figure 2D),	(Seq ID no 41)
40	CCG ACT TCA CCG GGT (selected from position 1334 in Figure 2D),	(Seq ID no 42)

MTC group (5S)

CGG AGG GGC AGT ATC (selected from positions 86-90 in Figure 3),

(Seq ID no 43)

Mycobacteria other than those of the MTC group (23S)

- GAT CAA TGC TCG GTT (selected from positions 99-101 in Figure 4A), (Seq ID no 44)
- 5 TTC CCC GCG TTA CCT (selected from position 183 in Figure 4A), (Seq ID no 45)
- TTA GCC TGT TCC GGT (selected from positions 261-271 in Figure 4A), (Seq ID no 46)
- GCA TGC GGT TTA GCC (selected from positions 281-284 in Figure 4B), (Seq ID no 47)
- TAC CCG GTT GTC CAT (selected from positions 290-293 in Figure 4B), (Seq ID no 48)
- GTA GAG CTG AGA CAT (selected from positions 327-335 and
- 10 343-357 in Figure 4B), (Seq ID no 49)
- GCC GTC CCA GGC CAC (selected from positions 400-405 in
- Figure 4B and Figure 4C), (Seq ID no 50)
- CTC GGG TGT TGA TAT (selected from positions 453-462 in Figure 4C), (Seq ID no 51)
- ACT ATT TCA CTC CCT (selected from positions 587-599 in Figure 4C), (Seq ID no 52)
- 15 ACG CCA TCA CCC CAC (selected from positions 637-660 in Figure 4D), (Seq ID no 53)
- CGA CGT GTC CCT GAC (selected from positions 704-712 in Figure 4D), (Seq ID no 54)
- ACT ACA CCC CAA AGG (selected from positions 763-789 in Figure 4E), (Seq ID no 55)
- CAC GCT TTT ACA CCA (selected from positions 1060-1074 in Figure 4E), (Seq ID no 56)
- GCG ACT ACA CAT CCT (selected from positions 1177-1185 in Figure 4E), (Seq ID no 57)
- 20 CGG CGC ATA ATC ACT (selected from positions 1259-1265 in Figure 4E), (Seq ID no 58)
- CCA CAT CCA CCG TAA (selected from positions 1311-1327 in Figure 4F), (Seq ID no 59)
- CGC TGA ATG GGG GAC (selected from positions 1345-1348 in Figure 4F), (Seq ID no 60)
- GGA GCT TCG CTG AAT (selected from positions 1361-1364 in Figure 4G), (Seq ID no 61)
- CGG TCA CCC GGA GCT (selected from positions 1361-1364 in Figure 4G), (Seq ID no 62)
- 25 GGA CGC CCA TAC ACG (selected from positions 1556-1570 in Figure 4G), (Seq ID no 63)
- GAA GGG GAA TGG TCG (selected from positions 1608-1613 in Figure 4H), (Seq ID no 64)
- AAT CGC CAC GCC CCC (selected from positions 1626-1638 in Figure 4H), (Seq ID no 65)
- CAG CGA AGG TCC CAC (selected from positions 1651-1659 in Figure 4H), (Seq ID no 66)
- GTC ACC CCA TTG CTT (selected from positions 1675-1677 in Figure 4H), (Seq ID no 67)
- 30 ATC GCT CTC TAC GGG (selected from positions 1734-1741 in Figure 4H), (Seq ID no 68)
- GTG TAT GTG CTC GCT (selected from positions 1847-1853 in Figure 4I), (Seq ID no 69)
- ACG GTA TTC CGG GCC (selected from positions 1967-1976 in Figure 4I), (Seq ID no 70)
- GGC CGA ATC CCG CTC (selected from positions 2006-2010 in Figure 4I), (Seq ID no 71)
- AAA CAG TCG CTA CCC (selected from positions 2025-2027 in Figure 4I), (Seq ID no 72)
- 35 CCT TAC GGG TTA ACG (selected from positions 2131-2132 in Figure 4J), (Seq ID no 73)
- GAG ACA GTT GGG AAG (selected from positions 2252-2255 in Figure 4J), (Seq ID no 74)
- TGG CGT CTG TGC TTC (selected from positions 2396-2405 in
- Figure 4J and Figure 4K), (Seq ID no 75)
- CGA CTC CAC ACA AAC (selected from positions 2416-2420 in Figure 4K), (Seq ID no 76)
- 40 GAT AAG GGT TCG ACG (selected from positions 2474-2478 in Figure 4K), (Seq ID no 77)
- ATC CGT TGA GTG ACA (selected from position 2687 in Figure 4K), (Seq ID no 78)
- CAG CCC GTT ATC CCC (selected from position 2719 in Figure 4K), (Seq ID no 79)

- AAC CTT TGG GAC CTG (selected from position 2809 in Figure 4L), (Seq ID no 80)
TAA AAG GGT GAG AAA (selected from positions 3062-3068 in Figure 4L), (Seq ID no 81)
GTC TGG CCT ATC AAT (selected from positions 3097-3106 in Figure 4L), (Seq ID no 82)
- 5 *Mycobacteria other than those of the MTC group (16S)*
AGA TTG CCC ACG TGT (selected from positions 135-136 in Figure 5A), (Seq ID no 83)
AAT CCG AGA AAA CCC (selected from positions 472-475 in Figure 5A), (Seq ID no 84)
GCA TTA CCC GCT GGC (selected from positions 1136-1144 in Figure 5A), (Seq ID no 85)
TTA AAA GGA TTC GCT (selected from positions 1287-1292 in Figure 5B), (Seq ID no 86)
10 AGA CCC CAA TCC GAA (selected from position 1313 in Figure 5B), (Seq ID no 87)
GAC TCC GAC TTC ATG (selected from position 1334 in Figure 5B), (Seq ID no 88)
- Drug resistance*
23S-mediated macrolide resistance (M. avium)
- 15 GTC TTT TCG TCC TGC (wild-type) (selected from positions 2568-2569 in Figure 6), (Seq ID no 89)
GTC TTA TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 90)
GTC TTC TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 91)
GTC TTG TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 92)
20 GTC TAT TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 93)
GTC TCT TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 94)
GTC TGT TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 95)
- 16S-mediated streptomycin resistance (M. tuberculosis)*
- 25 TTG GCC GGT GCT TCT (wild-type) (selected from positions 452 in Figure 7), (Seq ID no 96)
TTG GCC GGT ACT TCT (selected from positions 452 in Figure 7), (Seq ID no 97)
TTG GCC GGT CCT TCT (selected from positions 452 in Figure 7), (Seq ID no 98)
TTG GCC GGT TCT TCT (selected from positions 452 in Figure 7), (Seq ID no 99)
ACC GCG GCT GCT GGC (wild-type) (selected from positions 473-477 in Figure 7), (Seq ID no 100)
30 ACC GCG GCT ACT GGC (selected from positions 473 in Figure 7), (Seq ID no 101)
ACC GCG GCT CCT GGC (selected from positions 473 in Figure 7), or (Seq ID no 102)
ACC GCG GCT TCT GGC (selected from positions 473 in Figure 7), (Seq ID no 103)
CGG CAG CTG GCA CGT (selected from positions 474 in Figure 7), (Seq ID no 104)
35 CGG CCG CTG GCA CGT (selected from positions 474 in Figure 7), (Seq ID no 105)
CGG CTG CTG GCA CGT (selected from positions 474 in Figure 7), (Seq ID no 106)
CGT ATT ACC GCA GCT (selected from positions 477 in Figure 7), (Seq ID no 107)
CGT ATT ACC GCC GCT (selected from positions 477 in Figure 7), (Seq ID no 108)
CGT ATT ACC GCT GCT (selected from positions 477 in Figure 7), (Seq ID no 109)
40 TTC CTT TGA GTT TTA (wild-type) (selected from positions 865-866 in Figure 7), (Seq ID no 110)
TTC CTT TAA GTT TTA (selected from positions 865 in Figure 7), (Seq ID no 111)
TTC CTT TCA GTT TTA (selected from positions 865 in Figure 7), (Seq ID no 112)

TTC CTT TTA GTT TTA (selected from positions 865 in Figure 7), (Seq ID no 113)
 TTC CTT AGA GTT TTA (selected from positions 866 in Figure 7), (Seq ID no 114)
 TTC CTT CGA GTT TTA (selected from positions 866 in Figure 7), (Seq ID no 115)
 TTC CTT GGA GTT TTA (selected from positions 866 in Figure 7), (Seq ID no 116)

5

Other examples of suitable Q subsequences are given below.

CAT GTG TCC TGT GGT and (Seq ID no 117)
 CGT CAG CCC GAG AAA (Seq ID no 118)

10 selected so as to be complementary to *M. gordonae* 16S rRNA (positions 174-188 and 452-466, respectively, of GenBank entry GB:MSGRR16SI, accession no. M29563). These positions correspond to positions 192-206 and 473-487, respectively, of the alignments shown in Figure 2 and 5. Probes having this or a similar nucleobase sequence are suitable for detecting *M. gordonae*.

15

CAC TAC ACA CGC TCG, and (Seq ID no 119)
 TGG CGT TGA GGT TTC (Seq ID no 120)

selected so as to be complementary to positions 781-795 and 2369-2383, respectively, of *M. kansasii* 23S rRNA (GenBank entry MK23SRRNA accession number Z17212). These
 20 positions correspond to positions 774-794 and 2398-2412, respectively, of the alignments shown in Figure 1 and 4. Probes having this or a similar nucleobase sequence are suitable for detecting *M. kansasii*.

Precursor rRNA

25 AAC ACT CCC TTT GGA (Seq ID no 123)

A peptide nucleic acid probe having the above-indicated nucleobase sequence is directed to *M. tuberculosis* precursor rRNA. The probe is complementary to positions 602 to 616 of GenBank accession number X58890.

30

Especially, probes based on those nucleobase sequences with sequence identification numbers Seq ID no 62, 79 and 80 (and other probes selected from positions 1361-1364 in Figure 1F, 2719 in Figure 4K and 2809 in Figure 4L) are suitable for detecting *M. avium*. Probes based on the nucleobase sequence with sequence identification number Seq ID no 55
 35 (and other probes selected from positions 763-789 in Figure 4E) are suitable for detecting *M. avium*, *M. intracellulare* and *M. scrofulaceum* as a group (the organisms termed the MAIS group of mycobacteria). In addition, probes based on the nucleobase sequences with sequence identification numbers Seq ID no 77 and 81 are suitable for detecting *M. avium*, *M. intracellulare* and *M. paratuberculosis* as a group.

The invention is further illustrated by the non-limiting examples given below.

EXAMPLES

5

EXAMPLE 1

Mycobacterium species (*M. bovis* and *M. intracellulare*) 23S rDNA were partly amplified by PCR, and the PCR products were sequenced (both strands) using Cy5-labelled
10 oligonucleotide primers (DNA Technology, Aarhus, Denmark) and the 7-deaza-dGTP Thermo Sequenase cycle sequencing kit from Amersham, Little Chalfont, England. Sequences were read using an ALFexpress automated sequencer and ALFwin (version 1.10) software from Pharmacia Biotech, Uppsala, Sweden. *M. bovis* and *M. intracellulare* 23S rRNA sequences are included at the following positions of the 23S rDNA sequence alignments: positions 681-
15 729 (Figures 1C and 4D), positions 761-800 (Figures 1D and 4E), positions 2401-2440 (Figures 1H and 4K), positions 2441-2480 (Figures 1I and 4K), positions 2481-2520 (Figure 1I), positions 3041-3080 (Figure 4L), and positions 3081-3120 (Figures 1J and 4L).

EXAMPLE 2

20

Sequence alignments (see Figures 1 to 5) of 23S, 16S and 5S rDNA of mycobacteria of the MTC group, and 23S and 16S rDNA of mycobacteria other than those of the MTC group (MOTT) were done using the Megalign (version 3.12) alignment tool from DNASTAR (Madison, WI, USA). Up to one hundred sequences were aligned at a time.

25

Peptide nucleic acid probes in which the nucleobase sequence was complementary to distinctive mycobacterial rRNA were designed with due regard to secondary structures using the PrimerSelect program (version 3.04) from DNASTAR. As a control of sequence specificity, all probe sequences were subsequently matched with the GenBank and EMBL databases
30 using BLAST sequence similarity searching at the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov>).

As examples, the following sequences were selected:

35

MTC 23S

TCA CCA CCC TCC TCC

(Seq ID no 6)

CCA CCC TCC TCC

(modified Seq ID no 6)

ACT ATT CAC ACG CGC

(Seq ID no 8)

CCA CAC CCA CCA CAA

(Seq ID no 12)

	AAC TCC ACA CCC CCG	(Seq ID no 16)
	ACT CCA CAC CCC CGA	(Seq ID no 17)
	ACT CCG CCC CAA CTG	(Seq ID no 22)
	CTG TCC CTA AAC CCG	(Seq ID no 23)
5	TTC GAG GTT AGA TGC	(Seq ID no 24)
	GTC CCT AAA CCC GAT	(Seq ID no 25)
	GAC CTA TTG AAC CCG	(Seq ID no 29)

MTC 16S

10	GCA TCC CGT GGT CCT	(Seq ID no 33)
	CAC AAG ACA TGC ATC	(Seq ID no 34)
	GGC TTT TAA GGA TTC	(Seq ID no 40)

MOTT 23S

15	GAT CAA TGC TCG GTT	(Seq ID no 44)
	CGA CTC CAC ACA AAC	(Seq ID no 76)

MOTT 16S

	GCA TTA CCC GCT GGC	(Seq ID no 85)
20		

Drug resistance

	GTC TTA TCG TCC TGC	(Seq ID no 90)
	GTC TTC TCG TCC TGC	(Seq ID no 91)
	GTC TTG TCG TCC TGC	(Seq ID no 92)
25	GTC TAT TCG TCC TGC	(Seq ID no 93)
	GTC TCT TCG TCC TGC	(Seq ID no 94)
	GTC TGT TCG TCC TGC	(Seq ID no 95)

Precursor rRNA

30	AAC ACT CCC TTT GGA	(Seq ID no 123)
----	---------------------	-----------------

Non-sense probes

	GTC CGT GAA CCC GAT	(Seq ID no 121)
	TAC GCT CTT TGA GCT	(Seq ID no 122)
35		

EXAMPLE 3

Peptide nucleic acid probes were synthesised using an Expedite 8909 Nucleic Acid Synthesis System purchased from PerSeptive Biosystems (Framingham, USA). The peptide nucleic acid probes were terminated with two β -alanine molecules or with one or two lysine molecule(s) and, before cleavage from the resin, labelled with 5-(or 6)-carboxyfluorescein (Flu) or

rhodamine (Rho) at the β -amino group of alanine (peptide label) or ϵ -amino group of lysine (peptide label), respectively. Probes were purified using reverse phase HPLC at 50°C and characterised using a G2025 A MALDI-TOF MS instrument (Hewlett Packard, San Fernando, California, USA). Molecular weights determined were within 0.1% of the calculated molecular weights.

The following labelled peptide nucleic acid probes were synthesised:

MTC 23S

10	Lys(Flu)-Lys(Flu)-TCA CCA CCC TCC TCC-NH ₂	(OK 446/modified Seq ID no 6)
	Lys(Flu)-Lys(Flu)-CCA CCC TCC TCC-NH ₂	(OK 575/modified Seq ID no 6)
	Lys(Flu)-Lys(Flu)-ACT ATT CAC ACG CGC-NH ₂	(OK 447/modified Seq ID no 8)
	Lys(Flu)-ACT ATT CAC ACG CGC-NH ₂	(OK 688/modified Seq ID no 8)
	Lys(Flu)-Lys(Flu)-CCA CAC CCA CCA CAA-NH ₂	(OK 448/modified Seq ID no 12)
15	Lys(Flu)-Lys(Flu)-AAC TCC ACA CCC CCG-NH ₂	(OK 449/modified Seq ID no 16)
	Lys(Flu)-Lys(Flu)-ACT CCA CAC CCC CGA-NH ₂	(OK 309/modified Seq ID no 17)
	Lys(Flu)-Lys(Flu)-ACT CCG CCC CAA CTG-NH ₂	(OK 450/modified Seq ID no 22)
	Lys(Flu)-Lys(Flu)-CTG TCC CTA AAC CCG-NH ₂	(OK 305/modified Seq ID no 23)
	Lys(Flu)-Lys(Flu)-TTC GAG GTT AGA TGC-NH ₂	(OK 306/modified Seq ID no 24)
20	Lys(Flu)-TTC GAG GTT AGA TGC-NH ₂	(OK 682/modified Seq ID no 24)
	Lys(Flu)-Lys(Flu)-GTC CCT AAA CCC GAT-NH ₂	(OK 307/modified Seq ID no 25)
	Lys(Flu)-GTC CCT AAA CCC GAT-NH ₂	(OK 654/modified Seq ID no 25)
	Lys(Flu)-GAC CTA TTG AAC CCG-NH ₂	(OK 660/modified Seq ID no 29)

25 MTC 16S

	Lys(Flu)-Lys(Flu)-Gly-GCA TCC CGT GGT CCT-NH ₂	(OK 223/modified Seq ID no 33)
	Lys(Flu)-Lys(Flu)-CAC AAG ACA TGC ATC-NH ₂	(OK 310/modified Seq ID no 34)
	Lys(Flu)-CAC AAG ACA TGC ATC-NH ₂	(OK 655/modified Seq ID no 34)
	Lys(Flu)-GGC TTT TAA GGA TTC-NH ₂	(OK 689/modified Seq ID no 40)
30	Lys(Rho)-GGC TTT TAA GGA TTC-NH ₂	(OK 702/modified Seq ID no 40)

MOTT 23S

	Flu- β -Ala- β -Ala-GAT CAA TGC TCG GTT-NH ₂	(OK 624/modified Seq ID no 44)
	Flu- β -Ala- β -Ala-CGA CTC CAC ACA AAC-NH ₂	(OK 612/modified Seq ID no 76)

35

MOTT 16S

	Flu- β -Ala- β -Ala-GCA TTA CCC GCT GGC-NH ₂	(OK 623/modified Seq ID no 85)
--	---	--------------------------------

Drug resistance

40	Lys(Flu)-GTC TTT TCG TCC TGC-NH ₂	(OK 745/modified Seq ID no 89)
	Lys(Rho)-GTC TTA TCG TCC TGC-NH ₂	(OK 746/modified Seq ID no 90)

	Lys(Rho)-GTC TTC TCG TCC TGC-NH ₂	(OK 746/modified Seq ID no 91)
	Lys(Rho)-GTC TTG TCG TCC TGC-NH ₂	(OK 746/modified Seq ID no 92)
	Lys(Rho)-GTC TAT TCG TCC TGC-NH ₂	(OK 747/modified Seq ID no 93)
	Lys(Rho)-GTC TCT TCG TCC TGC-NH ₂	(OK 747/modified Seq ID no 94)
5	Lys(Rho)-GTC TGT TCG TCC TGC-NH ₂	(OK 747/modified Seq ID no 95)

Precursor rRNA

	Lys(Flu)-AAC ACT CCC TTT GGA-NH ₂	(OK 749/modified Seq ID no 123)
--	--	---------------------------------

10 Reduction of non-specific binding

	GTC CGT GAA CCC GAT-NH ₂	(OK 507/modified Seq ID no 121)
	Gly-TAC GCT CTT TGA GCT-NH ₂	(OK 714/modified Seq ID no 122)

EXAMPLE 4

15

Initially the ability of the peptide nucleic acid probes to react with target sequences of mycobacterial rRNA was tested by dot blot carried out with rRNA from *M. bovis* BCG, *M. avium* and *E. coli*.

20 *M. bovis* BCG (Statens Serum Institut, Denmark) and *M. intracellulare* (kindly provided by Statens Serum Institut) were grown in Dubos broth (Statens Serum Institut) or on Löwenstein-Jensen slants (Statens Serum Institut) at 37 °C. RNA was isolated from the bacterial cells using TRI-reagent (Sigma) following manufacture's directions. *E. coli* rRNA was purchased from Boehringer Mannheim, Germany.

25

200 ng *M. bovis* RNA, *M. intracellulare* RNA and *E. coli* rRNA were dotted onto membranes (Schleicher & Schüel, NY 13 N), and the membranes were dried and fixed under UV light for 2 minutes.

30 Protocol for dot blot assay

Each of the probes (70 nM probe in hybridisation solution (50 mM Tris, 10 mM NaCl, 10% (w/v) Dextran sulphate, 50% (v/v) glycerol, 5 mM EDTA, 0.1% (w/v) sodium pyrophosphate, 0.2% (w/v) polyvinylpyrrolidone, 0.2% (w/v) Ficoll, pH 7.6.)) were spotted onto a membrane. Hybridisation was continued for 1.5 hours at 55 or 65 °C, respectively. The membranes were
 35 rinsed 2 times for 15 minutes in 2 × SSPE buffer (1 × SSPE: 0.15 M NaCl, 10 mM sodium phosphate, 1 mM EDTA, pH 7.4) containing 0.1% SDS at ambient temperature, and subsequently 2 times for 15 minutes in 0.1 × SSPE buffer containing 0.1% SDS at 55 or 65 °C (see Table 1). The membrane was blocked with 0.5% (w/v) casein dissolved in 0.5M NaCl, 0.05M Tris/HCl pH 9.0. Thereafter, the membranes were incubated for 1 hour with rabbit-anti

FITC antibody labelled with alkaline phosphatase (AP) (DAKO K0046 vial A) diluted 1:2000 in 0.5% casein dissolved in 0.5M NaCl, 0.05M Tris/HCl pH 9.0. After incubation, the membranes were washed 3 times 5 minutes with TST buffer (0.05M Tris, 0.5M NaCl, 0.5% (w/v) Tween 20®, pH 9) at ambient temperature. Bound probes were visualised following standard
 5 procedures using BCIP/NBT, and the visualisation was stopped by incubation for 10 minutes with 10 mM EDTA. The blot was dried at 50 °C.

The results are given in Table 1 below.

10 TABLE 1

	E. coli rRNA		M. bovis BCG RNA		M. intracellulare RNA	
Probe	55 °C	65 °C	55 °C	65 °C	55 °C	65 °C
OK 305	negative	negative	positive	positive	negative	weak
OK 307	negative	negative	positive	positive	negative	weak
OK 309	negative	negative	positive	positive	negative	weak
OK 223	negative	negative	positive	positive	nd	nd
OK 310	negative	negative	negative	positive	negative	negative

nd: Not determined

The results indicate that all five peptide nucleic acid probes are capable of hybridising to target sequence of M. bovis BCG rRNA (as a representative of the MTC group), whereas no
 15 hybridisation to E. coli rRNA (as a representative of organisms other than mycobacteria) and no detectable hybridisation to M. intracellulare rRNA were observed (as a representative of the MOTT group).

EXAMPLE 5

20

This example illustrates the ability of the peptide nucleic acid probes to penetrate the mycobacterial cell wall and subsequently hybridise to target sequence of mycobacteria of the MTC group and not mycobacteria of the MOTT group, in particular not mycobacteria of the MAC group, or Neisseria gonorrhoeae, by fluorescence *in situ* hybridisation (FISH).

25

Preparation of bacterial slides

M. bovis BCG (Statens Seruminstitut, Denmark), M. avium (kindly provided by Statens Seruminstitut, Denmark), and M. intracellulare (kindly provided by Statens Seruminstitut,

Denmark) were grown in Dubos broth (Statens Seruminstitut, Denmark) or on Löwenstein-Jensen slants (Statens Seruminstitut, Denmark) at 37 °C. *N. gonorrhoeae* (Statens Seruminstitut, Denmark) was grown on chocolate agar (Statens Seruminstitut, Denmark) at 37 °C with additional 5% CO₂.

5

Cultures were smeared onto microscope slides and fixed according to standard procedures. Prior to the hybridisation, the smears were immersed into 80% ethanol for 15 minutes, and subsequently rinsed with water and air dried. This step is not essential for the following hybridisation step, but it is anticipated that it will kill any viable mycobacteria on the slides, and may further serve as an additional fixation step.

10

Protocol for fluorescence in situ hybridisation (FISH)

1. The bacterial slide was covered with a hybridisation solution containing the probe in question.
- 15 2. The slide was incubated in a humid incubation chamber at 45°C or 55°C for 90 minutes.
3. The slide was washed 25 minutes at 45°C or 55°C in prewarmed wash solution (5 mM Tris, 145 mM NaCl, pH 10) followed by 30 seconds in water.
4. The slide was dried and mounted with IMAGEN Mounting Fluid (DAKO, Copenhagen, Denmark)
- 20

The hybridisation solution contains 50 mM Tris, 10 mM NaCl, 10% (w/v) Dextran sulphate, 30% (v/v) formamide, 0.1% (v/v) Triton X-100®, 5 mM EDTA, 0.1% (w/v) sodium pyrophosphate, 0.2% (w/v) polyvinylpyrrolidone, 0.2% (w/v) Ficoll, pH 7.6.

25

Whenever possible, the applied equipment was heat-treated, and solutions were exposed to 1µl/ml diethylpyrocarbonate (Sigma Chemical Co.) in order to inactivate nucleases.

Microscopically examinations were conducted using a fluorescence microscope (Leica, Wetzlar, Germany) equipped with a 100x/1.20 water objective, a HBO 100 W lamp and a FITC filter set. Mycobacteria were identified as fluorescent, 1 - 10 µm slender, rod-shaped bacilli.

30

Fluorescein-labelled peptide nucleic acid probes targeting 23S rRNA of the mycobacteria of the MTC group (OK 306, OK 309, OK 446, OK 449) and 16S rRNA of the mycobacteria of the MTC group (OK 310) were tested. Individual probe concentrations and incubation temperatures are listed together with the results in Table 2 and 3.

35

TABLE 2

	OK 306 250nM 45°C	OK 309 250nM 45°C	OK 446 500nM 55°C	OK 449 500nM 55°C
M. bovis BCG	positive	positive	positive	positive
M. avium	negative	negative	negative	negative
M. intracellulare	negative	negative	not determined	not determined
N. gonorrhoeae	negative	negative	not determined	not determined

TABLE 3

	OK 447 1µM 55°C	OK 310 250nM 45°C	OK 306/OK 310 500/500nM 55°C
M. bovis BCG	positive	positive	positive
M. avium	negative	negative	negative
M. intracellulare	not determined	negative	negative
N. gonorrhoeae	not determined	negative	not determined

- 5 It can be concluded that the probes are able to penetrate the mycobacterial cell wall of mycobacterium cultures and subsequently hybridise to target rRNA sequence. This makes possible the development of fluorescence in situ hybridisation (FISH) protocols for specific detection of mycobacteria.

10 EXAMPLE 6

Test of probes on clinical smears of sputum

- The ability of the peptide nucleic acid to penetrate the cell wall of mycobacteria of the MTC group in clinical samples was tested on smears of sputum from suspected cases of tuberculosis (kindly provided by Division of Microbiology, Ramathibodi Hospital, Bangkok, Thailand) by fluorescence in situ hybridisation (FISH). Smears from the same patient were initially evaluated positive by Ziehl-Neelsen staining, which shows only the presence of acid fast bacilli, not whether these are mycobacteria of the MTC group.

- 20 Fluorescein-labelled peptide nucleic acid probes targeting 23S rRNA of the mycobacteria of the MTC group (OK 306, OK 446, OK 449) and 16S rRNA of the mycobacteria of the MTC group (OK 310) were used. Furthermore, a random peptide nucleic acid probe (a 15-mer wherein each position may be A, T, C or G (obtained from Millipore Corporation, Bedford, MA, USA) was added to the hybridisation solution in order to increase the signal-to-noise ratio.

FISH was carried out at 55 °C as described in Example 5. Applied probe concentrations are listed together with the results in Table 4 and 5.

TABLE 4

Sample number	OK 446/Random 1µM/50µM	OK 449/Random 1µM/50µM	Ziehl-Neelsen staining
285	Positive	Positive	4+
335	Positive	Eq.	2+
345	Positive	Positive	3+
224	Positive	Positive	3+
297	Negative	Eq.	2+
179	Negative	Negative	4+
247	Negative	Negative	2+
255	Positive	Positive	2+
202	Eq.	Positive	2+

5

TABLE 5

Sample number	OK 306/OK 310 500/500 nM	Ziehl-Neelsen staining
213	Positive	4+
292	Positive	4+
159	Positive	3+
287	Positive	3+

Smears stained by Ziehl-Neelsen staining were examined with a 100x objective and scored according to the following method: -: 0 bacilli, +/-: 1-200 per 300 fields, 2+: 1-9 per 10 fields, 3+: 1-9 per field, 4+: >9 per field.

- 10 Positive: Several mycobacteria were identified in the smear. Negative: No fluorescent mycobacteria were identified in the smear. Eq: Few (1-3) fluorescent mycobacteria were identified in the smear.

15 It appears from the table that the peptide nucleic acid probes are able to penetrate and subsequently hybridise to target sequence of mycobacteria of the MTC-group in AFB-positive sputum smears. The fact that not all AFB-positive sputum smears are found positive with applied probes indicate that not all AFB-positive sputum smears contains mycobacteria of the MTC-group.

EXAMPLE 7

20

The reactivity and specificity of selected peptide nucleic acid probes for detecting

mycobacteria of the MTC group as well as probes for detecting mycobacteria of the MOTT group were evaluated by fluorescence in situ hybridisation (FISH) on control smears prepared from cultures of different mycobacterium species. The mycobacterium species were selected so as to be representative for the mycobacterium genus as well as to include clinically relevant species.

M. tuberculosis (ATCC 25177), M. bovis BCG (ATCC 35734), M. intracellulare (ATCC 13950), M. avium (ATCC 25292), M. kansasii (ATCC12479), M. gordonae (ATCC 14470), M. scrofulaceum (ATCC 19981), M. abscessus (ATCC19977), M. marinum (ATCC 927), M. simiae (ATCC 25575), M. szulgai (ATCC 35799), M. flavescens (ATCC 23033), M. fortuitum (ATCC 43266) and M. xenopi (ATCC19250) were grown at Dubos broth (Statens Serum Institut) at 37 °C with the exception of M. marinum which was grown at 32 °C.

Smears were prepared as described in Example 5. FISH was carried out as described below.

Protocol for fluorescence in situ hybridisation (FISH)

1. The bacterial slide was covered with a hybridisation solution containing the probe in question.
2. The slide was incubated in a humid incubation chamber at 55°C for 90 minutes.
- 20 3. The slide was washed 30 minutes at 55°C in prewarmed wash solution (5 mM Tris, 15 mM NaCl, 0.1% (v/v), Triton X-100®, pH 10) followed by 30 seconds in water.
4. The slide was dried and mounted with IMAGEN Mounting Fluid (DAKO, Copenhagen, Denmark)
- 25 The hybridisation solution contained 50 mM Tris, 10 mM NaCl, 10% (w/v) Dextran sulphate, 30% (v/v) formamide, 0.1% (v/v) Triton X-100®, 5 mM EDTA, 0.1% (w/v) sodium pyrophosphate, 0.2% (w/v) polyvinylpyrrolidone, and 0.2% (w/v) Ficoll, pH 7.6. To avoid non-specific binding of the labelled peptide nucleic acid probe, 1-5 µM of non-labelled, non-sense peptide nucleic acid probe was added to the hybridisation solution (OK 507/modified Seq ID no 121 and/or OK 714/modified Seq ID no 122).
- 30

Whenever possible, the applied equipment was heat-treated, and solutions were exposed to 1µl/ml diethylpyrocarbonate (Sigma Chemical Co.) in order to inactivate nucleases.

35 Microscopic examinations were conducted using a fluorescence microscope (Leica, Wetzlar, Germany) equipped with a 100×/1.30 oil objective, a HBO 100 W lamp and a FITC/TRITC dual band filter set. Mycobacteria were identified on basis of both fluorescence (strong, medium, weak, no) and morphology (1-10 µm slender, rod-shaped bacilli. Mycobacteria of the MOTT

group may appear pleomorphic, ranging in appearance from long rods to coccoid forms)

Probe concentrations are listed together with the results in Table 6 and 7 (probes targeting mycobacteria of the MTC group) and Table 8 (probes targeting to mycobacteria of the MOTT group).

5

TABLE 6

	OK 450 25 nM	OK 682 100 nM	OK 689 100 nM	OK 688 250 nM	OK 660 100 nM
<i>M. tuberculosis</i>	+++	+++	+++	+++	+++
<i>M. bovis</i> BCG	+++	+++	+++	+++	+++
<i>M. intracellulare</i>	-	-	-	-	-
<i>M. avium</i>	-	-	-	-	-
<i>M. kansasii</i>	++	-	-	-	-
<i>M. gordonae</i>	-	-	-	-	-
<i>M. scrofulaceum</i>	+++	-	-	-	-
<i>M. abscessus</i>	-	-	-	-	+
<i>M. marinum</i>	+++	-	+	+	+++
<i>M. simiae</i>	-	-	-	-	-
<i>M. szulgai</i>	+++	-	-	-	-
<i>M. flavescens</i>	-	++	-	-	-
<i>M. fortuitum</i>	-	+	-	-	-
<i>M. xenopi</i>	-	++	-	-	-

+++ strong fluorescence, ++ medium fluorescence, + weak fluorescence, - no fluorescence

TABLE 7

Mycobacteria	OK 655 150 nM	OK 448 50 nM	OK 654 100 nM	OK 446 25 nM
<i>M. tuberculosis</i>	+++	+++	+++	+++
<i>M. bovis</i> BCG	+++	+++	+++	+++
<i>M. intracellulare</i>	-	-	-	-
<i>M. avium</i>	-	-	-	-
<i>M. kansasii</i>	-	-	-	-
<i>M. gordonae</i>	-	-	-	-
<i>M. scrofulaceum</i>	-	-	-	-
<i>M. abscessus</i>	-	-	+	-
<i>M. marinum</i>	-	-	+	+++
<i>M. simiae</i>	-	-	-	-
<i>M. szulgai</i>	-	-	-	-
<i>M. flavescens</i>	-	-	-	-
<i>M. fortuitum</i>	-	-	-	-
<i>M. xenopi</i>	-	-	-	-

+++ strong fluorescence, ++ medium fluorescence, + weak fluorescence, - no fluorescence

TABLE 8

Mycobacteria	OK 612 100 nM	OK 624 100 nM	OK 623 100 nM
<i>M. tuberculosis</i>	-	-	-
<i>M. bovis</i> BCG	-	-	-
<i>M. intracellulare</i>	-	++	++
<i>M. avium</i>	+++	+++	+++
<i>M. kansasii</i>	-	-	+++
<i>M. gordonae</i>	-	++	++
<i>M. scrofulaceum</i>	-	++	++
<i>M. abscessus</i>	-	++	+++
<i>M. marinum</i>	-	-	-
<i>M. simiae</i>	-	++	+++
<i>M. szulgai</i>	-	-	+++
<i>M. flavescens</i>	-	-	-
<i>M. fortuitum</i>	-	++	-
<i>M. xenopi</i>	-	-	-

5 +++ strong fluorescence, ++ medium fluorescence, + weak fluorescence, - no fluorescence

Each of probes indicated in Table 6, 7 and 8 was further investigated with regard to hybridisation to other common respiratory bacteria, namely *Corynebacterium* spp.,

Fusobacterium nucleatum, *Haemophilus influenzae*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Propionibacterium acnes*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Brahmella catarrhalis*, *Escherichia coli*, *Neisseria* spp., *Actinobacter calcoaceticus*, *Actinomyces* spp., *Enterobacter aerogenes*, *Proteus mirabilis*, *Pseudomonas maltophilia*,
5 *Streptococcus viridans*, and *Nocardia asteroides*. No cross-hybridisation was observed by fluorescence in situ hybridisation to any of these bacteria in the case of OK 682, OK 654, OK 655, OK 688, OK 660, OK 612, OK 624 and OK 623. Some cross-reactivity was observed in the case of OK 446 (to *P. acnes*), OK 448 (to *P. acnes* and *B. catarrhalis*), and OK 450 (to *P. acnes* and *B. catarrhalis*).

10

Table 6 and 7 shows that none of the MTC probes cross-react with *M. intracellulare* and/or *M. avium*, but indeed strongly with *M. tuberculosis* and *M. bovis* BCG. As shown in Table 8, both OK 624 and OK 623 hybridise to *M. intracellulare* and *M. avium* which are both members of the MAC group, whereas none of them hybridise to *M. tuberculosis* or *M. bovis* BCG. OK 612
15 hybridises to *M. avium* only. It should be noted that the aligned sequence of *M. intracellulare* has just one nucleobase difference to the target sequence of *M. avium*, see Figure 4K.

The data support the use of the methodology described in claim 3 and 4 and exemplified in Example 2 for design of peptide nucleic acid probes that are capable of hybridising to target
20 sequence of one or more mycobacterium species and not to other mycobacterium species having at least one nucleobase difference to the target sequence.

EXAMPLE 8

25 To study the usefulness of the peptide nucleic acid probes in distinguishing between mycobacteria of the MTC group and mycobacteria of the MOTT group, the probes were tested on smears of mycobacterium-positive cultures prepared from 34 + 28 clinical samples (sputum samples, other respiratory samples and extrapulmonary samples) from individuals suspected of tuberculosis or other mycobacterial infections (kindly provided by the Mycobacterium
30 Department, Statens Serum Institut, Denmark). Complex/species identification data obtained with the AccuProbe tests from Gen-Probe Inc., USA were available for each sample.

Table 9 shows the results obtained with four different peptide nucleic acid probes targeting mycobacteria of the MTC group (OK 682, OK 660, OK 688 and OK 689) and one probe
35 targeting mycobacteria of the MOTT group (OK 623), and Table 10 shows the results obtained with two peptide nucleic acid probes targeting mycobacteria of the MOTT group (OK 623 and OK 612) and a mixture of two probes targeting mycobacteria of the MTC group (OK 688 and OK 689). Data are arranged according to the results obtained by AccuProbe. Sample

preparation, hybridisation and visualisation were performed as described in Example 7.

TABLE 9

Complex/ species (n)	OK 623 25 nM n_p	OK 682 100 nM n_p	OK 660 100 nM n_p	OK 688 250 nM n_p	OK 689 100 nM n_p
MTC (23)	0	23	23	23	23
<i>M. avium</i> (5)	5	0	0	0	0
<i>M. gordonae</i> (3)	3	0	0	0	0
Unknown (3)	3	0	0	0	0

n_p denotes number of positive samples.

- 5 * The term "unknown" means that the sample not contains mycobacteria of the MTC group, or mycobacteria of the MAC group according the AccuProbe test, but further species identification was not performed.

TABLE 10

Complex/ species (n)	OK 623 25nM n_p	OK 612 100 nM n_p	OK 688/OK 689 50 nM/50 nM n_p
MTC (17)	0		16
<i>M. avium</i> (2)	2	2	0
<i>M. gordonae</i> (4)	3	0	0
Unknown (5)	5	0	0

- 10 n_p denotes number of positive samples.

* The term "unknown" means that the sample not contains mycobacteria of the MTC group, or mycobacteria of the MAC group according to the AccuProbe test, but further species identification was not performed.

- 15 The results shown in Table 9 are in conformity with the complex/species identification performed with the AccuProbe tests, and thus confirm that peptide nucleic acid probes can be used to determine whether an infection is caused by mycobacteria of the MTC group or by mycobacteria of the MOTT group.
- 20 From the results in Table 10, it can be seen that it is possible to differentiate between mycobacteria of the MTC group and mycobacteria of the MOTT group with 100% specificity and 91-94% sensitivity relative to results obtained by the AccuProbe tests. Furthermore, OK 612 is very suitable for specific identification of *M. avium* among those being positive for mycobacteria of the MOTT group as the result is positive in the case of *M. avium* and negative
- 25 in the other cases of mycobacteria of the MOTT group.

EXAMPLE 9

Direct detection of mycobacteria in clinical smears of sputum

This example demonstrates the ability of the peptide nucleic acid to detect and identify

- 5 mycobacteria directly in AFB-positive sputum samples from suspected cases of tuberculosis (kindly provided by Division of Microbiology, Ramathibodi Hospital, Bangkok, Thailand) and suspected cases of other mycobacterial infections (kindly provided by Clinical Microbiology Dept., Rigshospitalet, Copenhagen, Denmark) by FISH is shown.

- 10 The clinical smears were prepared according to the procedure described in Example 5, and FISH was performed as described in Example 7. The results are shown in Table 11.

TABLE 11

Sample no.	OK 623 25 nM	OK 654 100 nM	OK 655 150 nM	OK 682 100 nM	OK 688 250 nM	OK 689 100 nM
1	-	++	++	++	++	++
175	-	++	nd	nd	++	++
459	-	-	nd	nd	-	-
166	-	-	-	nd	-	-
268	-	++	++	++	++	++
34267	++	-	-	-	-	-

nd: not determined

- 15 +++ strong fluorescence, ++ medium fluorescence, + weak fluorescence, - no fluorescence

- It appears from examples in Table 11 that AFB-positive sputum smears were evaluated positive for mycobacteria of the MTC group (sample numbers 1, 175, and 268), positive for mycobacteria of the MOTT group (sample number 37267), or negative for mycobacteria (sample numbers. 459 and 166) by the applied probes. Thus, PNA-probes are useful reagents for specific identification of mycobacteria directly in sputum smears by fluorescence in situ hybridisation. AFB-positive sputum samples that are negative with all probes may be explained in three ways: a) the sample may contain mycobacteria not detected by the probes, e.g. *M. fortuitum*, b) the sample may contain other acid-fast bacteria than mycobacteria, or c) the mycobacteria in the sample lack or have a strongly reduced content of rRNA due to for example antibiotic treatment.
- 20
- 25

- In conclusion, direct identification of mycobacteria in smear-positive sputum samples by peptide nucleic acid-based fluorescence in situ hybridisation combines simplicity and morphological advantages of current staining methods with concomitant species identification, and will thus allow clinical microbiology laboratories to benefit from the
- 30

advantages offered by molecular techniques to provide crucial information pertaining to therapy and patient management.

EXAMPLE 10

5

This example demonstrates simultaneous detection and identification of mycobacteria of the MTC group and mycobacteria of the MOTT group using differently labelled probes targeting mycobacteria of the MTC group and mycobacteria of the MOTT group, respectively, by fluorescence in situ hybridisation.

10

Control smears of different mycobacterium species were prepared as described in Example 5. In addition, smears containing a mixture of *M. tuberculosis* and *M. avium* were prepared (Table 8, last row). FISH was performed as described in Example 7.

15

A rhodamine-labelled peptide nucleic acid probe targeting 16S rRNA of mycobacteria of the MTC group (OK 702) and a fluorescein-labelled peptide nucleic acid probe targeting 16S rRNA of mycobacteria of the MOTT group (OK 623) were applied simultaneously in the concentrations listed in Table 12 together with the results.

20

TABLE 12

Mycobacterium species	OK 623/OK 702 25/250 nM
<i>M. tuberculosis</i>	- (G)/ +++ (R)
<i>M. bovis</i> BCG	- (G)/ +++ (R)
<i>M. avium</i>	+++ (G)/ - (R)
<i>M. intracellulare</i>	+++ (G)/ - (R)
<i>M. kansasii</i>	+++ (G)/ - (R)
<i>M. avium</i> / <i>M. tuberculosis</i>	+++ (G)/+++ (R)

+++ strong fluorescence - no fluorescence

G green fluorescence, R red fluorescence

Mycobacteria of the MTC group, i.e. *M. tuberculosis* and *M. bovis*, were observed as green fluorescent mycobacteria, whereas mycobacteria of the MOTT group, i.e. *M. avium*, *M. intracellulare* and *M. kansasii*, were observed as red fluorescent mycobacteria. Mycobacteria in the *M. avium*/*M. tuberculosis* mixture were identified by a mixture of both green fluorescent mycobacteria and red fluorescent mycobacteria.

The results show that it is possible to distinguish between different *Mycobacterium* species in

one smear using a mixture of differently labelled probes. Such simultaneous detection and identification of mycobacteria may further be extended to comprise three or more differently labelled peptide nucleic acid probes.

5 EXAMPLE 11

The ability of a peptide nucleic acid probes to hybridise to precursor rRNA and further to distinguish between precursor rRNA of *M. tuberculosis* and precursor rRNA of *M. avium* was investigated by fluorescence in situ hybridisation.

10

Smears were prepared as described in Example 5 and FISH were carried out as described in Example 7 using a fluorescein-labelled probe targeting precursor rRNA of *M. tuberculosis* (OK 749). The results are given in Table 13.

15 TABLE 13

Mycobacterium	OK 749 1000 nM
<i>M. tuberculosis</i>	+
<i>M. avium</i>	-

+ weak fluorescence - no fluorescence

From the results, it can be concluded that it is possible to detect precursor rRNA, and further that it is possible to distinguish between precursor rRNA from different mycobacterium species.

20 The application of peptide nucleic acid targeting precursor rRNA may be particularly useful for measuring the mycobacterial growth and thus be an indicator of the viability of the mycobacteria. This would in particular be important for monitoring of the effect of antibiotics in relation to both treatment of tuberculosis and drug susceptibility studies.

25 EXAMPLE 12

The ability of peptide nucleic acid probes for differentiation of drug susceptible and drug resistant mycobacteria was evaluated using a fluorescein-labelled probe targeting the wild type sequence of 23S rRNA of *M. avium* and *M. intracellulare* together with rhodamine-labelled probes targeting single point mutations associated with macrolide resistance in *M.*
30 *avium* and *M. intracellulare*.

Smears were prepared as described in Example 5 from cultures of *M. avium* (ATCC no. 25292) and *M. intracellulare* (ATCC no. 13950). These strains are anticipated to contain the

wild type sequence of rRNA. Macrolide resistant variants were not available. FISH was carried out as described in Example 7 using a fluorescein-labelled peptide nucleic acid probe targeting wild type 23S rRNA (OK 745) and a mixture of rhodamine-labelled peptide nucleic acid probes targeting the three possible mutations at position 2568 (OK 746) and at position 52569 (OK 747) of *M. avium* 23S rDNA of GenBank entry X52917 (see Figure 6). The results are given in Table 14.

TABLE 14

Mycobacterium species	OK 745/OK 746/OK 747 500/500/500 nM
<i>M. avium</i> (wild type)	+++ (G)/ - (R)
<i>M. intracellulare</i> (wild type)	+++ (G)/ - (R)

+++ strong fluorescence - no fluorescence

10 G green fluorescence, R red fluorescence

OK 746 and OK 747 are each a mixture of three single point mutation probes

The results in Table 14 show that *M. avium* and *M. intracellulare* are detected with the fluorescein-labelled probe (OK 745) targeting *M. avium* and *M. intracellulare* wild types and not detected with the mixture of rhodamine-labelled probes (OK 746 and OK 747) targeting single point mutations associated with macrolide resistance. Such peptide nucleic acid probes targeting the wild type and drug resistant variants, respectively, may be important tools for both the prediction of an efficient therapy as well as for monitoring the effect of the treatment.

20

EXAMPLE 13

To illustrate the speed with which peptide nucleic acid probes penetrate the mycobacterial cell wall and subsequently hybridise to their target sequence the protocol described in Example 7 was modified to 15 minutes hybridisation time and the results compared with 90 minutes hybridisation time. Smears were prepared as described in Example 5. The results are given in Table 15.

25

TABLE 15

	OK 623 25 nM		OK 689 100 nM	
	15 min	90 min	15 min	90 min
M. tuberculosis			++	+++
M. avium	++	+++		

+++ strong fluorescence ++ medium fluorescence
+ weak fluorescence - no fluorescence

- 5 The data presented in Table 15 show that hybridisation by peptide nucleic acid probes inside the mycobacterial cells is accomplished in a very short time resulting in a detectable signal after just 15 minutes incubation. Thus, the use peptide nucleic acid probes makes possible the development of very fast fluorescence in situ hybridisation protocols.

10 EXAMPLE 14

To describe the ability of very short peptide nucleic acid probes to hybridise to target sequences, a 12-mer peptide nucleic acid probe labelled with fluorescein (OK 575) was tested by fluorescence in situ hybridisation (FISH).

15

Smears were prepared as described in Example 5 and FISH were carried out as described in Example 7. The results are given in Table 16.

TABLE 16

Mycobacterium	OK 575 50 nM
M. tuberculosis	+
M. bovis BCG	++
M. avium	-
M. intracellulare	-
M. kansasii	-

20 ++ medium fluorescence + weak fluorescence - no fluorescence

The results in table 17 shows that a 12-mer peptide nucleic acid probe is capable of hybridising specifically to target sequences under the same stringency conditions as 15-mers.

A lower fluorescence intensity is obtained as the T_m for a 12-mer peptide nucleic acid probe is

25 lower than T_m for a 15-mer peptide nucleic acid probe.

The data clearly suggest that by lowering the stringency condition, e.g. by decreasing the hybridisation/washing temperature and/or the concentration of formamide, even shorter probes may be applied for detection of mycobacteria provided that specific sequences of such can be designed.

CLAIMS

1. Peptide nucleic acid probe for detecting a target sequence of one or more mycobacteria optionally present in a sample, said probe being capable of hybridising to a target sequence of mycobacterial rDNA, precursor rRNA or rRNA forming detectable hybrids, and a mixture of such probes.
2. Peptide nucleic acid probe according to claim 1, said probe being capable of hybridising to a target sequence of mycobacterial rDNA, precursor rRNA, or 23S, 16S or 5S rRNA forming detectable hybrids, and a mixture of such probes.
3. Peptide nucleic acid probe according to claim 1 or 2, said probe being capable of hybridising to a target sequence of mycobacterial rDNA, precursor rRNA, or 23S, 16S or 5S rRNA forming detectable hybrids, said target sequence being obtainable by
- (a) comparing the nucleobase sequences of said mycobacterial rRNA or rDNA of one or more mycobacteria to be detected with the corresponding nucleobase sequence of organism(s), in particular other mycobacteria, in particular other mycobacteria, from which said one or more mycobacteria are to be distinguished,
- (b) selecting a target sequence of said rRNA or rDNA which includes at least one nucleobase differing from the corresponding nucleobase of the organism(s), in particular other mycobacteria, from which said one or more mycobacteria are to be distinguished, and
- (c) determining the capability of said probe to hybridise to the selected target sequence to form detectable hybrids, and a mixture of such probes.
4. Peptide nucleic acid probe according to claim 1 or 2, said probe being capable of hybridising to a target sequence of mycobacterial rDNA, precursor rRNA or 23S, 16S or 5S rRNA forming detectable hybrids, said probe being obtainable by
- (a) comparing the nucleobase sequences of said mycobacterial rRNA or rDNA of one or more mycobacteria to be detected with the corresponding nucleobase sequence of organism(s), in particular other mycobacteria, in particular other mycobacteria, from which said one or more mycobacteria are to be distinguished,

(b) selecting a target sequence of said rRNA or rDNA which includes at least one nucleobase differing from the corresponding nucleobase of the organism(s), in particular other mycobacteria, from which said one or more mycobacteria are to be distinguished,


5 (c) synthesising said probe, and

(d) determining the capability of said probe to hybridise to the selected target sequence to form detectable hybrids,
and a mixture of such probes.

10

5. Peptide nucleic acid probe according to any one of claims 1 to 4 for detecting a target sequence of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC) or for detecting a target sequence of one or more mycobacteria other than mycobacteria of the Mycobacterium tuberculosis Complex (MOTT) optionally present in a sample, which probe
15 comprises from 6 to 30 polymerised peptide nucleic acid moieties, said probe being capable of hybridising to a target sequence of mycobacterial rDNA, precursor rRNA or 23S, 16S or 5S rRNA forming detectable hybrids,
and a mixture of such probes.

20 6. Peptide nucleic acid probe according to any one of claims 1 to 5 for detecting a target sequence of rDNA, precursor rRNA or 23S, 16S or 5S rRNA of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC) or for detecting a target sequence of rDNA, precursor rRNA or 23S, 16S or 5S rRNA of one or more mycobacteria other than mycobacteria of the Mycobacterium tuberculosis Complex (MOTT) optionally present in a
25 sample, which probe comprises from 10 to 30 polymerised moieties of formula (I)



(I)

30

wherein each X and Y independently designate O or S,
each Z independently designates O, S, NR¹, or C(R¹)₂, wherein each R¹ independently designate H, C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl,
35 each R², R³ and R⁴ designate independently H, the side chain of a naturally occurring amino acid, the side chain of a non-naturally occurring amino acid, C₁₋₄ alkyl, C₁₋₄ alkenyl or C₁₋₄ alkynyl, or a functional group, each Q independently designates a naturally occurring nucleobase, a non-naturally occurring nucleobase, an intercalator, a nucleobase-binding

group, a label or H,

with the proviso that the probe comprising such subsequence is capable of forming detectable hybrids with the target sequence of said mycobacterial rDNA, precursor rRNA or 23S, 16S or

5 5S rRNA,

and a mixture of such probes.

7. Peptide nucleic acid probe according to any one of claims 1 to 6 for detecting a target sequence of 23S rRNA of one or more mycobacteria of the *Mycobacterium tuberculosis*

10 Complex (MTC) optionally present in a sample, which probe comprises from 10 to 30 polymerised moieties of formula (I) as defined in claim 6,

with the proviso that the Qs of adjacent moieties are selected so as to form a sequence of which a subsequence includes at least one nucleobase that is complementary to a

15 nucleobase of *M. tuberculosis* 23S rRNA differing from the corresponding nucleobase of at least *M. avium* located within the following domains

Positions 149-158 in Figure 1A,

Positions 220-221 in Figure 1A,

20 Positions 328-361 in Figure 1A and Figure 1B,

Positions 453-455 in Figure 1B,

Positions 490-501 in Figure 1B,

Positions 637-660 in Figure 1C,

Positions 706-712 in Figure 1D,

25 Positions 762-789 in Figure 1D,

Position 989 in Figure 1D,

Positions 1068-1072 in Figure 1D,

Position 1148 in Figure 1E,

Positions 1311-1329 in Figure 1E,

30 Positions 1361-1364 in Figure 1F,

Position 1418 in Figure 1F,

Positions 1563-1570 in Figure 1F,

Positions 1627-1638 in Figure 1G,

Positions 1675-1677 in Figure 1G,

35 Position 1718 in Figure 1G,

Positions 1734-1740 in Figure 1H,

Positions 1967-1976 in Figure 1H,

Positions 2403-2420 in Figure 1H,

Positions 2457-2488 in Figure 1I,
Positions 2952-2956 in Figure 1I,
Positions 2966-2969 in Figure 1J,
Positions 3000-3003 in Figure 1J or
5 Positions 3097-3106 in Figure 1J,

and further with the proviso that the probe comprising such subsequence is capable of forming detectable hybrids with a target sequence of said mycobacterial 23S rRNA, and a mixture of such probes.

10

8. Peptide nucleic acid probe according to any one of claims 1 to 6 for detecting a target sequence of 16S rRNA of one or more mycobacteria of the *Mycobacterium tuberculosis* Complex (MTC) optionally present in a sample, which probe comprises from 10 to 30 polymerised moieties of formula (I) as defined in claim 6,

15

with the proviso that the Qs of adjacent moieties are selected so as to form a sequence of which a subsequence includes at least one nucleobase that is complementary to a nucleobase of *M. tuberculosis* 16S rRNA differing from the corresponding nucleobase of at least *M. avium* located within the following domains

20

Positions 76-79 in Figure 2A,
Positions 98-101 in Figure 2A,
Positions 135-136 in Figure 2A,
Positions 194-201 in Figure 2B,
25 Positions 222-229 in Figure 2B,
Position 242 in Figure 2B,
Position 474 in Figure 2C,
Positions 1136-1145 in Figure 2C,
Positions 1271-1272 in Figure 2C,
30 Positions 1287-1292 in Figure 2D,
Position 1313 in Figure 2D, or
Position 1334 in Figure 2D,

and further with the proviso that the probe comprising such subsequence is capable of forming detectable hybrids with a target sequence of said mycobacterial 16S rRNA, and a mixture of such probes.

35

9. Peptide nucleic acid probe according to any one of claims 1 to 6 for detecting a target

sequence of 5S rRNA of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC) optionally present in a sample, which probe comprises from 10 to 30 polymerised moieties of formula (I) as defined in claim 6,

- 5 with the proviso that the Qs of adjacent moieties are selected so as to form a sequence of which a subsequence includes at least one nucleobase that is complementary to a nucleobase of M. tuberculosis 5S rRNA differing from the corresponding nucleobase of at least M. avium located within the following domain

- 10 Positions 86-90 in Figure 3

and further with the proviso that the probe comprising such subsequence is capable of forming detectable hybrids with a target sequence of said mycobacterial 5S rRNA, and a mixture of such probes.

15

10. Peptide nucleic acid probe according to any one of claims 1 to 8 for detecting a target sequence of 23S or 16S rRNA of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC) optionally present in a sample, which probe comprises from 10 to 30 polymerised moieties of formula (I) as defined in claim 6,

20

with the proviso that the Qs of adjacent moieties are selected so as to form a sequence of which a subsequence includes at least one nucleobase that is complementary to a nucleobase of M. tuberculosis 23S or 16 S rRNA differing from the corresponding nucleobase of at least M. avium located within the following domains

25

Positions 149-158 in Figure 1A,
Positions 328-361 in Figure 1A and Figure 1B,
Positions 490-501 in Figure 1B,
Positions 637-660 in Figure 1C,

30

Positions 762-789 in Figure 1D,
Positions 1068-1072 in Figure 1D,
Positions 1311-1329 in Figure 1E,
Positions 1361-1364 in Figure 1F,
Positions 1563-1570 in Figure 1F,

35

Positions 1627-1638 in Figure 1G,
Positions 1734-1740 in Figure 1H,
Positions 2457-2488 in Figure 1I,
Positions 2952-2956 in Figure 1I,

Positions 3097-3106 in Figure 1J,
Positions 135-136 in Figure 2 A, or
Positions 1287-1292 in Figure 2D,

- 5 and further with the proviso that the probe comprising such subsequence is capable of forming detectable hybrids with a target sequence of said mycobacterial 23S or 16S rRNA, and a mixture of such probes.

11. Peptide nucleic acid probe according to any one of claims 1 to 6 for detecting a target
10 sequence of 23S rRNA of one or more mycobacteria other than mycobacteria of the *Mycobacterium tuberculosis* Complex (MOTT) optionally present in a sample, which probe comprises from 10 to 30 polymerised moieties of formula (I) as defined in claim 6,

- with the proviso that the Qs of adjacent moieties are selected so as to form a sequence of
15 which a subsequence includes at least one nucleobase that is complementary to a nucleobase of *M. avium* 23S rRNA differing from the corresponding nucleobase of at least *M. tuberculosis* located within the following domains

- Positions 99-101 in Figure 4A,
20 Position 183 in Figure 4A,
Positions 261-271 in Figure 4A,
Positions 281-284 in Figure 4B,
Positions 290-293 in Figure 4B,
Positions 327-335 in Figure 4B,
25 Positions 343-357 in Figure 4B,
Positions 400-405 in Figure 4B and Figure 4C,
Positions 453-462 in Figure 4C,
Positions 587-599 in Figure 4C,
Positions 637-660 in Figure 4D,
30 Positions 704-712 in Figure 4D,
Positions 763-789 in Figure 4E,
Positions 1060-1074 in Figure 4E,
Positions 1177-1185 in Figure 4E,
Positions 1259-1265 in Figure 4F,
35 Positions 1311-1327 in Figure 4F,
Positions 1345-1348 in Figure 4F,
Positions 1361-1364 in Figure 4G,
Positions 1556-1570 in Figure 4G,

- Positions 1608-1613 in Figure 4H,
Positions 1626-1638 in Figure 4H,
Positions 1651-1659 in Figure 4H,
Positions 1675-1677 in Figure 4H,
5 Positions 1734-1741 in Figure 4H,
Positions 1847-1853 in Figure 4I,
Positions 1967-1976 in Figure 4I,
Positions 2006-2010 in Figure 4I,
Positions 2025-2027 in Figure 4I,
10 Positions 2131-2132 in Figure 4J,
Positions 2252-2255 in Figure 4J,
Positions 2396-2405 in Figure 4J and Figure 4K,
Positions 2416-2420 in Figure 4K,
Positions 2474-2478 in Figure 4K,
15 Position 2687 in Figure 4K,
Position 2719 in Figure 4K,
Position 2809 in Figure 4L,
Positions 3062-2068 in Figure 4L, or
Positions 3097-3106 in Figure 4L,
20
- and further with the proviso that the probe comprising such subsequence is capable of forming detectable hybrids with a target sequence of said mycobacterial 23S rRNA, and a mixture of such probes.
- 25 12. Peptide nucleic acid probe according to any one of claims 1 to 6 for detecting a target sequence of 16S rRNA of one or more mycobacteria other than mycobacteria of the *Mycobacterium tuberculosis* Complex (MOTT) optionally present in a sample, which probe comprises from 10 to 30 polymerised moieties of formula (I) as defined in claim 6,
- 30 with the proviso that the Qs of adjacent moieties are selected so as to form a sequence of which a subsequence includes at least one nucleobase that is complementary to a nucleobase of *M. avium* 16S rRNA differing from the corresponding nucleobase of at least *M. tuberculosis* located within the following domains
- 35 Positions 135-136 in Figure 5A,
Positions 472-475 in Figure 5A,
Positions 1136-1144 in Figure 5A,
Positions 1287-1292 in Figure 5B,

Position 1313 in Figure 5B, or
Position 1334 in Figure 5B,

and further with the proviso that the probe comprising such subsequence is capable of forming
5 detectable hybrids with a target sequence of said mycobacterial 16S rRNA,
and a mixture of such probes.

13. Peptide nucleic acid probe according to any one of claims 1 to 6, 11 and 12 for detecting a
target sequence of 23S or 16S rRNA of one or more mycobacteria other than mycobacteria of
10 the Mycobacterium tuberculosis Complex (MOTT) optionally present in a sample, which probe
comprises from 10 to 30 polymerised moieties of formula (I) as defined in claim 6,

with the proviso that the Qs of adjacent moieties are selected so as to form a sequence of
which a subsequence includes at least one nucleobase that is complementary to a
15 nucleobase of *M. avium* 23S or 16S rRNA differing from the corresponding nucleobase of at
least *M. tuberculosis* located within the following domains

Positions 99-101 in Figure 4A,
Positions 290-293 in Figure 4B,
20 Positions 400-405 in Figure 4B and Figure 4C,
Positions 453-462 in Figure 4C,
Positions 637-660 in Figure 4D,
Positions 763-789 in Figure 4E,
Positions 1311-1327 in Figure 4F,
25 Positions 1361-1364 in Figure 4G,
Positions 1734-1741 in Figure 4H,
Positions 2025-2027 in Figure 4I,
Positions 2474-2478 in Figure 4K,
Positions 3062-2068 in Figure 4L, or
30 Positions 1287-1292 in Figure 5B,

and further with the proviso that the probe comprising such subsequence is capable of forming
detectable hybrids with a target sequence of said mycobacterial 23S or 16S rRNA,
and a mixture of such probes.

35

14. Peptide nucleic acid probe according to any one of claims 1 to 6 for detecting a target
sequence of 23S, 16S or 5S rRNA of one or more mycobacteria of the Mycobacterium
tuberculosis Complex (MTC) or for detecting a target sequence of 23S, 16S or 5S rRNA of

one or more mycobacteria other than mycobacteria of the Mycobacterium tuberculosis Complex (MOTT) optionally present in a sample, which probe comprises from 10 to 30 polymerised moieties of formula (I) as defined in claim 6,

- 5 with the proviso that the Qs of adjacent moieties are selected so as to form a sequence of which a subsequence includes at least one nucleobase that is complementary to a nucleobase that differs from the corresponding nucleobase of 23S, 16S or 5S rRNA of said one or more mycobacteria located within the following domains

10 positions 2568-2569 in Figure 6,

Position 452 in Figure 7,




Positions 473-477 in Figure 7, or

Positions 865-866 in Figure 7,

15

and further with the proviso that the probe comprising such subsequence is capable of forming detectable hybrids with the target sequence of said mycobacterial 23S, 16S or 5S rRNA, and a mixture of such probes.

- 20 15. Peptide nucleic acid probe according to any one of claims 1 to 14 of formula (II), (III), or (IV)



wherein Z, R², R³, and R⁴, and Q is as defined in claim 6 with the provisos defined in claims 6

to 14,
and a mixture of such probes.

16. Peptide nucleic acid probe according to any one of claims 1 to 15, wherein Z is NH, NCH₃
5 or O, each R², R³ and R⁴ independently designate H or the side chain of a naturally occurring amino acid, the side chain of a non-naturally occurring amino acid, or C₁₋₄ alkyl, and each Q is a naturally occurring nucleobase or a non-naturally occurring nucleobase with the provisos defined in claims 6 to 14,
and a mixture of such probes.


10

17. Peptide nucleic acid probe according to any one of claims 1 to 16, wherein Z is NH or O,
and R² is H or the side chain of Ala, Asp, Cys, Glu, His, HomoCys, Lys, Orn, Ser or Thr, and
Q is a nucleobase selected from thymine, adenine, cytosine, guanine, uracil, iso-C and 2,6-
diaminopurine with the provisos defined in claims 6 to 14,

15 and a mixture of such probes.

18. Peptide nucleic acid probe according to any one of claims 1 to 17 of formula (V)

20



(V)

wherein R⁴ is H or the side chain of Ala, Asp, Cys, Glu, His, HomoCys, Lys, Orn, Ser or Thr,
and Q is as defined in claim 17 with the provisos defined in claims 6 to 14,
25 and a mixture of such probes.

25

19. Peptide nucleic acid probe according to any one of claims 1 to 18 further comprising one
or more labels and a mixture of such probes, which labels may be mutually identical or
different, which probes optionally may comprise one or more linkers, and which probes may
30 be mutually identical or different with the provisos defined in claims 6 to 14.

30

20. Peptide nucleic acid probe according to any one of claims 1 to 19 for detecting a target
sequence of one or more mycobacteria, the nucleobase sequence of said probe being
substantially complementary to the nucleobase sequence of said target sequence.

35

21. Peptide nucleic acid probe according to any one of claims 1 to 20 for detecting a target
sequence of one or more mycobacteria, the nucleobase sequence of said probe being
complementary to the nucleobase sequence of said target sequence.

22. Peptide nucleic acid probes according to any one of claims 1 to 21, wherein the Qs of adjacent moieties are selected so as to form the following subsequences

- | | | |
|----|---|----------------|
| 5 | AGA TGC GGG TAG CAC (selected from positions 149-158 in Figure 1A), | (Seq ID no 1) |
| | TGT TTT CTC CTC CTA (selected from positions 220-221 in Figure 1A), | (Seq ID no 2) |
| | ACT GCC TCT CAG CCG (selected from positions 328-361 in | |
| | Figure 1A and Figure 1B), | (Seq ID no 3) |
| | TGA TAC TAG GCA GGT (selected from positions 453-455 in Figure 1B), | (Seq ID no 4) |
| 10 | CGG ATT CAC AGC GGA (selected from positions 490-501 in Figure 1B), | (Seq ID no 5) |
| | TCA CCA CCC TCC TCC (selected from positions 637-660 in Figure 1C), | (Seq ID no 6) |
| | TTA ACC TTG CGA CAT (selected from positions 706-712 in Figure 1C), | (Seq ID no 7) |
| | ACT ATT CAC ACG CGC (selected from positions 762-789 in Figure 1D), | (Seq ID no 8) |
| | CTC CGC GGT GAA CCA (selected from position 989 in Figure 1D), | (Seq ID no 9) |
| 15 | GCT TTA CAC CAC GGC (selected from positions 1068-1072 in Figure 1D), | (Seq ID no 10) |
| | ACG CTT GGG GGC CTT (selected from position 1148 in Figure 1E), | (Seq ID no 11) |
| | CCA CAC CCA CCA CAA (selected from positions 1311-1329 in Figure 1E), | (Seq ID no 12) |
| | CCG GTG GCT TCG CTG (selected from positions 1361-1364 in Figure 1F), | (Seq ID no 13) |
| | ACT TGC CTT GTC GCT (selected from position 1418 in Figure 1F), | (Seq ID no 14) |
| 20 | GAT TCG TCA CGG GCG (selected from positions 1563-1570 in Figure 1F), | (Seq ID no 15) |
| | AAC TCC ACA CCC CCG (selected from positions 1627-1638 in Figure 1G), | (Seq ID no 16) |
| | ACT CCA CAC CCC CGA (selected from positions 1627-1638 in Figure 1G), | (Seq ID no 17) |
| | ACC CCT TCG CTT GAC (selected from positions 1675-1677 in Figure 1G), | (Seq ID no 18) |
| | CTT GCC CCA GTG TTA (selected from position 1718 in Figure 1G), | (Seq ID no 19) |
| 25 | CTC TCC CTA CCG GCT (selected from positions 1734-1740 in Figure 1H), | (Seq ID no 20) |
| | GAT ATT CCG GTC CCC (selected from positions 1967-1976 in Figure 1H), | (Seq ID no 21) |
| | ACT CCG CCC CAA CTG (selected from positions 2403-2420 in Figure 1H), | (Seq ID no 22) |
| | CTG TCC CTA AAC CCG (selected from positions 2457-2488 in Figure 1I), | (Seq ID no 23) |
| | TTC GAG GTT AGA TGC (selected from positions 2457-2488 in Figure 1I), | (Seq ID no 24) |
| 30 | GTC CCT AAA CCC GAT (selected from positions 2457-2488 in Figure 1I), | (Seq ID no 25) |
| | GGT GCA CCA GAG GTT (selected from positions 2952-2956 in Figure 1I), | (Seq ID no 26) |
| | CTG GCG GGA CAA CTG (selected from positions 2966-2969 in Figure 1J), | (Seq ID no 27) |
| | TTA TCC TGA CCG AAC (selected from positions 3000-3003 in Figure 1J), | (Seq ID no 28) |
| | GAC CTA TTG AAC CCG (selected from positions 3097-3106 in Figure 1J), | (Seq ID no 29) |
| 35 | | |
| | GAA GAG ACC TTT CCG (selected from positions 76-79 in Figure 2A), | (Seq ID no 30) |
| | CAC TCG AGT ATC TCC (selected from positions 98-101 in Figure 2A), | (Seq ID no 31) |
| | ATC ACC CAC GTG TTA (selected from positions 136-136 in Figure 2A), | (Seq ID no 32) |
| | GCA TCC CGT GGT CCT (selected from positions 194-201 in Figure 2B), | (Seq ID no 33) |
| 40 | CAC AAG ACA TGC ATC (selected from positions 194-201 in Figure 2B), | (Seq ID no 34) |
| | TAA AGC GCT TTC CAC (selected from positions 222-229 in Figure 2B), | (Seq ID no 35) |
| | GCT CAT CCC ACA CCG (selected from position 242 in Figure 2B), | (Seq ID no 36) |

- CCG AGA GAA CCC GGA (selected from position 474 in Figure 2C), (Seq ID no 37)
 AGT CCC CAC CAT TAC (selected from positions 1136-1145 in Figure 2C), (Seq ID no 38)
 AAC CTC GCG GCA TCG (selected from positions 1271-1272 in Figure 2C), (Seq ID no 39)
 GGC TTT TAA GGA TTC (selected from positions 1287-1292 in Figure 2D), (Seq ID no 40)
 5 GAC CCC GAT CCG AAC (selected from position 1313 in Figure 2D), (Seq ID no 41)
 CCG ACT TCA CGG GGT (selected from position 1334 in Figure 2D), (Seq ID no 42)
- CGG AGG GGC AGT ATC (selected from positions 86-90 in Figure 3), (Seq ID no 43)
- 10 GAT CAA TGC TCG GTT (selected from positions 99-101 in Figure 4A), (Seq ID no 44)
 TTC CCC GCG TTA CCT (selected from position 183 in Figure 4A), (Seq ID no 45)
 TTA GCC TGT TCC GGT (selected from positions 261-271 in Figure 4A), (Seq ID no 46)
 GCA TGC GGT TTA GCC (selected from positions 281-284 in Figure 4B), (Seq ID no 47)
 TAC CCG GTT GTC CAT (selected from positions 290-293 in Figure 4B), (Seq ID no 48)
- 15 GTA GAG CTG AGA CAT (selected from positions 327-335 and
 343-357 in Figure 4B), (Seq ID no 49)
 GCC GTC CCA GGC CAC (selected from positions 400-405 in
 Figure 4B and Figure 4C), (Seq ID no 50)
 CTC GGG TGT TGA TAT (selected from positions 453-462 in Figure 4C), (Seq ID no 51)
- 20 ACT ATT TCA CTC CCT (selected from positions 587-599 in Figure 4C), (Seq ID no 52)
 ACG CCA TCA CCC CAC (selected from positions 637-660 in Figure 4D), (Seq ID no 53)
 CGA CGT GTC CCT GAC (selected from positions 704-712 in Figure 4D), (Seq ID no 54)
 ACT ACA CCC CAA AGG (selected from positions 763-789 in Figure 4E), (Seq ID no 55)
 CAC GCT TTT ACA CCA (selected from positions 1060-1074 in Figure 4E), (Seq ID no 56)
- 25 GCG ACT ACA CAT CCT (selected from positions 1177-1185 in Figure 4E), (Seq ID no 57)
 CGG CGC ATA ATC ACT (selected from positions 1259-1265 in Figure 4F), (Seq ID no 58)
 CCA CAT CCA CCG TAA (selected from positions 1311-1327 in Figure 4F), (Seq ID no 59)
 CGC TGA ATG GGG GAC (selected from positions 1345-1348 in Figure 4F), (Seq ID no 60)
 GGA GCT TCG CTG AAT (selected from positions 1361-1364 in Figure 4G), (Seq ID no 61)
- 30 CGG TCA CCC GGA GCT (selected from positions 1361-1364 in Figure 4G), (Seq ID no 62)
 GGA CGC CCA TAC ACG (selected from positions 1556-1570 in Figure 4G), (Seq ID no 63)
 GAA GGG GAA TGG TCG (selected from positions 1608-1613 in Figure 4H), (Seq ID no 64)
 AAT CGC CAC GCC CCC (selected from positions 1626-1638 in Figure 4H), (Seq ID no 65)
 CAG CGA AGG TCC CAC (selected from positions 1651-1659 in Figure 4H), (Seq ID no 66)
- 35 GTC ACC CCA TTG CTT (selected from positions 1675-1677 in Figure 4H), (Seq ID no 67)
 ATC GCT CTC TAC GGG (selected from positions 1734-1741 in Figure 4H), (Seq ID no 68)
 GTG TAT GTG CTC GCT (selected from positions 1847-1853 in Figure 4I), (Seq ID no 69)
 ACG GTA TTC CGG GCC (selected from positions 1967-1976 in Figure 4I), (Seq ID no 70)
 GGC CGA ATC CCG CTC (selected from positions 2006-2010 in Figure 4I), (Seq ID no 71)
- 40 AAA CAG TCG CTA CCC (selected from positions 2025-2027 in Figure 4I), (Seq ID no 72)
 CCT TAC GGG TTA ACG (selected from positions 2131-2132 in Figure 4J), (Seq ID no 73)
 GAG ACA GTT GGG AAG (selected from positions 2252-2255 in Figure 4J), (Seq ID no 74)
 TGG CGT CTG TGC TTC (selected from positions 2396-2405 in

- Figure 4J and Figure 4K), (Seq ID no 75)
 CGA CTC CAC ACA AAC (selected from positions 2416-2420 in Figure 4K), (Seq ID no 76)
 GAT AAG GGT TCG ACG (selected from positions 2474-2478 in Figure 4K), (Seq ID no 77)
 ATC CGT TGA GTG ACA (selected from position 2687 in Figure 4K), (Seq ID no 78)
 5 CAG CCC GTT ATC CCC (selected from position 2719 in Figure 4K), (Seq ID no 79)
 AAC CTT TGG GAC CTG (selected from position 2809 in Figure 4L), (Seq ID no 80)
 TAA AAG GGT GAG AAA (selected from positions 3062-3068 in Figure 4L), (Seq ID no 81)
 GTC TGG CCT ATC AAT (selected from positions 3097-3106 in Figure 4L), (Seq ID no 82)
- 10 AGA TTG CCC ACG TGT (selected from positions 135-136 in Figure 5A), (Seq ID no 83)
 AAT CCG AGA AAA CCC (selected from positions 472-475 in Figure 5A), (Seq ID no 84)
 GCA TTA CCC GCT GGC (selected from positions 1136-1144 in Figure 5B), (Seq ID no 85)
 TTA AAA GGA TTC GCT (selected from positions 1287-1292 in Figure 5B), (Seq ID no 86)
 AGA CCC CAA TCC GAA (selected from position 1313 in Figure 5B), (Seq ID no 87)
 15 GAC TCC GAC TTC ATG (selected from position 1334 in Figure 5B), (Seq ID no 88)
- GTC TTT TCG TCC TGC (selected from positions 2568-2569 in Figure 6), (Seq ID no 89)
 GTC TTA TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 90)
 GTC TTC TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 91)
 20 GTC TTG TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 92)
 GTC TAT TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 93)
 GTC TCT TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 94)
 GTC TGT TCG TCC TGC (selected from positions 2568 in Figure 6), (Seq ID no 95)
- 25 TTG GCC GGT GCT TCT (selected from positions 452 in Figure 7), (Seq ID no 96)
 TTG GCC GGT ACT TCT (selected from positions 452 in Figure 7), (Seq ID no 97)
 TTG GCC GGT CCT TCT (selected from positions 452 in Figure 7), (Seq ID no 98)
 TTG GCC GGT TCT TCT (selected from positions 452 in Figure 7), (Seq ID no 99)
 ACC GCG GCT GCT GGC (selected from positions 473-477 in Figure 7), (Seq ID no 100)
 30 ACC GCG GCT ACT GGC (selected from positions 473 in Figure 7), (Seq ID no 101)
 ACC GCG GCT CCT GGC (selected from positions 473 in Figure 7), or (Seq ID no 102)
 ACC GCG GCT TCT GGC (selected from positions 473 in Figure 7), (Seq ID no 103)
 CGG CAG CTG GCA CGT (selected from positions 474 in Figure 7), (Seq ID no 104)
 CGG CCG CTG GCA CGT (selected from positions 474 in Figure 7), (Seq ID no 105)
 35 CGG CTG CTG GCA CGT (selected from positions 474 in Figure 7), (Seq ID no 106)
 CGT ATT ACC GCA GCT (selected from positions 477 in Figure 7), (Seq ID no 107)
 CGT ATT ACC GCC GCT (selected from positions 477 in Figure 7), (Seq ID no 107)
 CGT ATT ACC GCT GCT (selected from positions 477 in Figure 7), (Seq ID no 109)
 TTC CTT TGA GTT TTA (selected from positions 865-866 in Figure 7), (Seq ID no 110)
 40 TTC CTT TAA GTT TTA (selected from positions 865 in Figure 7), (Seq ID no 111)
 TTC CTT TCA GTT TTA (selected from positions 865 in Figure 7), (Seq ID no 112)
 TTC CTT TTA GTT TTA (selected from positions 865 in Figure 7), (Seq ID no 113)
 TTC CTT AGA GTT TTA (selected from positions 866 in Figure 7), (Seq ID no 114)

	TTC CTT CGA GTT TTA (selected from positions 866 in Figure 7),	(Seq ID no 115)
	TTC CTT GGA GTT TTA (selected from positions 866 in Figure 7),	(Seq ID no 116)
	CAT GTG TCC TGT GGT	(Seq ID no 117)
	CGT CAG CCC GAG AAA	(Seq ID no 118)
5	CAC TAC ACA CGC TCG	(Seq ID no 119)
	TGG CGT TGA GGT TTC and	(Seq ID no 120)
	AAC ACT CCC TTT GGA	(Seq ID no 123)

and a mixture of such probes.

10

23. Peptide nucleic acid probes according to claim 22, wherein the Qs of adjacent moieties are selected so as to form the following subsequences

	TCA CCA CCC TCC TCC	(Seq ID no 6)
15	CCA CCC TCC TCC	(modified Seq ID no 6)
	ACT ATT CAC ACG CGC	(Seq ID no 8)
	CCA CAC CCA CCA CAA	(Seq ID no 12)
	AAC TCC ACA CCC CCG	(Seq ID no 16)
	ACT CCA CAC CCC CGA	(Seq ID no 17)
20	ACT CCG CCC CAA CTG	(Seq ID no 22)
	CTG TCC CTA AAC CCG	(Seq ID no 23)
	TTC GAG GTT AGA TGC	(Seq ID no 24)
	GTC CCT AAA CCC GAT	(Seq ID no 25)
	GAC CTA TTG AAC CCG	(Seq ID no 29)
25	GCA TCC CGT GGT CCT	(Seq ID no 33)
	CAC AAG ACA TGC ATC	(Seq ID no 34)
	GGC TTT TAA GGA TTC	(Seq ID no 40)
30	GAT CAA TGC TCG GTT	(Seq ID no 44)
	CGA CTC CAC ACA AAC	(Seq ID no 76)
	GCA TTA CCC GCT GGC	(Seq ID no 85)
35	GTC TTA TCG TCC TGC	(Seq ID no 90)
	GTC TTC TCG TCC TGC	(Seq ID no 91)
	GTC TTG TCG TCC TGC	(Seq ID no 92)
	GTC TAT TCG TCC TGC	(Seq ID no 93)
	GTC TCT TCG TCC TGC	(Seq ID no 94)
40	GTC TGT TCG TCC TGC	(Seq ID no 95)
	AAC ACT CCC TTT GGA	(Seq ID no 123)

- CAT GTG TCC TGT GGT (Seq ID no 117)
 CGT CAG CCC GAG AAA (Seq ID no 118)
- 5 CAC TAC ACA CGC TCG, (Seq ID no 119)
 TGG CGT TGA GGT TTC (Seq ID no 120)

and a mixture of such probes.

- 10 24. Peptide nucleic acid probes according to claim 22 or 23 selected from

- Lys(Flu)-Lys(Flu)-TCA CCA CCC TCC TCC-NH₂ (OK 446/modified Seq ID no 6)
 Lys(Flu)-Lys(Flu)-CCA CCC TCC TCC-NH₂ (OK 575/modified Seq ID no 6)
 Lys(Flu)-Lys(Flu)-ACT ATT CAC ACG CGC-NH₂ (OK 447/modified Seq ID no 8)
 15 Lys(Flu)-ACT ATT CAC ACG CGC-NH₂ (OK 688/modified Seq ID no 8)
 Lys(Flu)-Lys(Flu)-CCA CAC CCA CCA CAA-NH₂ (OK 448/modified Seq ID no 12)
 Lys(Flu)-Lys(Flu)-AAC TCC ACA CCC CCG-NH₂ (OK 449/modified Seq ID no 16)
 Lys(Flu)-Lys(Flu)-ACT CCA CAC CCC CGA-NH₂ (OK 309/modified Seq ID no 17)
 Lys(Flu)-Lys(Flu)-ACT CCG CCC CAA CTG-NH₂ (OK 450/modified Seq ID no 22)
 20 Lys(Flu)-Lys(Flu)-CTG TCC CTA AAC CCG-NH₂ (OK 305/modified Seq ID no 23)
 Lys(Flu)-Lys(Flu)-TTC GAG GTT AGA TGC-NH₂ (OK 306/modified Seq ID no 24)
 Lys(Flu)-TTC GAG GTT AGA TGC-NH₂ (OK 682/modified Seq ID no 24)
 Lys(Flu)-Lys(Flu)-GTC CCT AAA CCC GAT-NH₂ (OK 307/modified Seq ID no 25)
 Lys(Flu)-GTC CCT AAA CCC GAT-NH₂ (OK 654/modified Seq ID no 25)
 25 Lys(Flu)-GAC CTA TTG AAC CCG-NH₂ (OK 660/modified Seq ID no 29)
- Lys(Flu)-Lys(Flu)-Gly-GCA TCC CGT GGT CCT-NH₂ (OK 223/modified Seq ID no 33)
 Lys(Flu)-Lys(Flu)-CAC AAG ACA TGC ATC-NH₂ (OK 310/modified Seq ID no 34)
 Lys(Flu)-CAC AAG ACA TGC ATC-NH₂ (OK 655/modified Seq ID no 34)
 30 Lys(Flu)-GGC TTT TAA GGA TTC-NH₂ (OK 689/modified Seq ID no 40)
 Lys(Rho)-GGC TTT TAA GGA TTC-NH₂ (OK 702/modified Seq ID no 40)
- Flu-β-Ala-β-Ala-GAT CAA TGC TCG GTT-NH₂ (OK 624/modified Seq ID no 44)
 Flu-β-Ala-β-Ala-CGA CTC CAC ACA AAC-NH₂ (OK 612/modified Seq ID no 76)
 35 Flu-β-Ala-β-Ala-GCA TTA CCC GCT GGC-NH₂ (OK 623/modified Seq ID no 85)
- Lys(Flu)-GTC TTT TCG TCC TGC-NH₂ (OK 745/modified Seq ID no 89)
 Lys(Rho)-GTC TTA TCG TCC TGC-NH₂ (OK 746/modified Seq ID no 90)
 40 Lys(Rho)-GTC TTC TCG TCC TGC-NH₂ (OK 746/modified Seq ID no 91)
 Lys(Rho)-GTC TTG TCG TCC TGC-NH₂ (OK 746/modified Seq ID no 92)
 Lys(Rho)-GTC TAT TCG TCC TGC-NH₂ (OK 747/modified Seq ID no 93)

Lys(Rho)-GTC TCT TCG TCC TGC-NH₂

(OK 747/modified Seq ID no 94)

Lys(Rho)-GTC TGT TCG TCC TGC-NH₂

(OK 747/modified Seq ID no 95)

Lys(Flu)-AAC ACT CCC TTT GGA-NH₂

(OK 749/modified Seq ID no 123)

5

wherein Flu denotes a 5-(and 6)-carboxyfluorescein label and Rho denotes a rhodamine label,

and a mixture of such probes.

10 25. Use of a peptide nucleic acid probe according to any one of claims 1 to 24 or a mixture thereof for detecting a target sequence of one or more mycobacteria optionally present in a sample.

26. Use of a peptide nucleic acid probe or a mixture thereof according to claim 25 for detecting
15 a target sequence of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC), in particular a target sequence of *M. tuberculosis*.

27. Use of a peptide nucleic acid probe or a mixture thereof according to claims 25 for detecting a target sequence of one or more mycobacteria other than mycobacteria of the
20 Mycobacterium tuberculosis Complex, in particular a target sequence of one or more mycobacteria of the Mycobacterium avium Complex.

28. Method for detecting a target sequence of one or more mycobacteria optionally present in a sample comprising

25

(1) contacting any rRNA or rDNA present in said sample with one or more peptide nucleic acid probes according to any one of claims 1 to 24 or a mixture thereof under conditions, whereby hybridisation takes place between said probe(s) and said rRNA or rDNA, and

30

(2) observing or measuring any formed detectable hybrids, and relating said observation or measurement to the presence of a target sequence of one or more mycobacteria in said sample.

35 29. Method according to claim 28 for detecting a target sequence of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC), in particular a target sequence of *M. tuberculosis*.

30. Method according to claim 28 for detecting a target sequence of one or more

mycobacteria other than mycobacteria of the Mycobacterium tuberculosis Complex.

31. Method according to any one of claims 28 to 30, wherein the hybridisation takes place in situ.

5

32. Method according to any of of claims 28 to 30, wherein the hybridisation takes place in vitro.

33. A method according to any one of claims 28 to 32,
10 c h a r a c t e r i s e d in that a signal amplifying system is used for measuring the resulting hybridisation.

34. Method according to any one of claims 28 to 33, wherein the sample is a sputum sample.

15 35. Kit for detecting a target sequence of one or more mycobacteria, in particular a target sequence of one or more mycobacteria of the Mycobacterium tuberculosis Complex (MTC), in particular a target sequence of M. tuberculosis, and/or for detecting a target sequence of one or more mycobacteria other than mycobacteria of the Mycobacterium tuberculosis Complex (MOTT), in particular a target sequence of one or more mycobacteria of the Mycobacterium
20 avium Complex,
c h a r a c t e r i s e d in that said kit comprises at least one peptide nucleic acid probe according to any one of claims 1 to 24, and optionally a detection system with at least one detecting reagent.

25 36. Kit according to claim 35,
c h a r a c t e r i s e d in that it further comprises a solid phase capture system.

1/31




Figure 1A

2/31

	370	380	390	400	
1327	CAGTCAGAAAGTGT	CGTGGTTAGCGGAAGTGGCCTGGGGAT			M.tuberculosis
656	TAGTCAGAAAGTGT	CGTGGTTAGCGGAAGTGGCCTGGGA			M.avium
656	TAGTCAGAAAGTGT	CGTGGTTAGCGGAAGTGGCCTGGGA			M.paratuberc.
742	TAGTGAATAAAGCAGT	GTGGTTAGGTGAAGTGGCCTGGGGAT			M.phlei
668	TAGTCAGAAAGTG	CGTGGTTAGCGGAATGGCCTGGGGAT			M.leprae
443	CAGTCAGAAAGTGT	CGTGGTTAGCGGAAGTGGCCTGGGGAT			M.gastri
386	CAGTCAGAAAGTGT	CGTGGTTAGCGGAAGTGGCCTGGGGAT			M.kansasii
2823	CAGTGAATAATGT	GTGGTTAGCGGAATGGCTTGGGAT			M.smegmatis

- - - -

	450	460	470	480	
1406	CGGCACCTGCCTAGT	TATCAATTCCCGAGTAGCAGCGGGCC			M.tuberculosis
735	CGGCACCTGCCTTAT	TATCAATCCCGAGTAGCAGCGGGCC			M.avium
735	CGGCACCTGCCTTAT	TATCAATCCCGAGTAGCAGCGGGCC			M.paratuberc.
820	TGCTGCCGCTGT	CACAGG--TCCCGAGTAGCAGCGGGCC			M.phlei
747	TGGCACCTGCCTTGT	TATCAATTCCCGAGTAGCAGCGGGCC			M.leprae
522	CGGCACCTGCCTTGT	TATCAATTCCCGAGTAGCAGCGGGCC			M.gastri
465	CGGCACCTGCCTTGT	TATCAATTCCCGAGTAGCAGCGGGCC			M.kansasii
2902	CGACCTCTGCTTGT	GATGGTGTCCCGAGTAGCAGCGGGCC			M.smegmatis

	490	500	510	520	
1446	CGTGGAATCGCTGTGAATCG	CCCGGGACCACCCGGTAAG			M.tuberculosis
775	CGTGGAATCGCTGTGAATCG	CCCGGGACCACCCGGTAAG			M.avium
775	CGTGGAATCGCTGTGAATCG	CCCGGGACCACCCGGTAAG			M.paratuberc.
857	CGTGGAATCGCTGTGAATCG	CCCGGGACCACCCGGTAAG			M.phlei
787	CGTGGAATCGCTGTGAATCG	CCCGGGACCACCCGGTAAG			M.leprae
562	CGTGGAATCGCTGTGAATCG	CCCGGGACCACCCGGTAAG			M.gastri
505	CGTGGAATCGCTGTGAATCG	CCCGGGACCACCCGGTAAG			M.kansasii
2942	CGTGGAATCGCTGTGAATCG	CCCGGGACCACCCGGTAAG			M.smegmatis

Figure 1B

3/31




Figure 1C

4/31

	770	780	790	800	
1726	CGACCCACACGCGCATACGCGCGTGTGAATAGTGGCGTGT				M.tuberculosis
84	CGACCCACACGCGCATACGCGCGTGTGAATAGTGGCGTGT				M.bovis
1039	CG	-----CATCCCTTTTGGGGTGT-----	AGTGGCGTGT		M.avium
103	CG	-----CATCCCTTTTGGGGTGT-----	AGTGGCGTGT		M.intracellula
1039	CG	-----CATCCCTTTTGGGGTGT-----	AGTGGCGTGT		M.paratuberc.
1126	CGTAT	CCACCTGTGTTGGGGTTGGTGT	AGTGGTGTGT		M.phlei
1052	CGTAT	CACGTTGTGAGCGTGTGT	AGTGGCGTGT		M.leprae
827	CGTAT	CACGCGTAAGCGTGTGT	AGTGGCGTGT		M.gastri
770	CGTAT	CGCGCGSAGCGTGTGT	AGTGGCGTGT		M.kansasii
3212	CGTAT	CCACACATAGTGTGTGT	AGTGGTGTGT		M.smegmatis

- - - -

	970	980	990	1000	
1926	ATTTAGGTGCAGCGTTGCGTGGTTCACCGCGGAGGTAGAG				M.tuberculosis
1228	ATTTAGGTGCAGCGTTGCGTGGTTCACCGCGGAGGTAGAG				M.avium
1228	ATTTAGGTGCAGCGTTGCGTGGTTCACCGCGGAGGTAGAG				M.paratuberc.
1322	ATTTAGGTGCAGCGT	CGCATGTTTCTTAT	CGGAGGTAGAG		M.phlei
1244	ATTTAGGTGCAGCGTTGCGTGGTTCACCGCGGAGGTAGAG				M.leprae
1019	ATTTAGGTGCAGCGTTGCGTGGTTCACCGCGGAGGTAGAG				M.gastri
962	ATTTAGGTGCAGCGTTGCGTGGTTCACCGCGGAGGTAGAG				M.kansasii
3408	ATTTAGGTGCAGCGT	CGCATGTTTCTTGC	CGGAGGTAGAG		M.smegmatis

- - - -

	1050	1060	1070	1080	
2005	CAGCCAAACTCCGAATGCCG-TGGTG-TA-AA-TCGTGGCA				M.tuberculosis
1307	CAGCCAAACTCCGAATGCCG-TGGTG-TAAAAGCGTGGCA				M.avium
1307	CAGCCAAACTCCGAATGCCG-TGGTG-TAAAAGCGTGGCA				M.paratuberc.
1401	CAGCCAAACTCCGAATGCCG	ATAAG-T	TAAGTGTGGCA		M.phlei
1323	CAGCCAAACTCCGAATGCCG-TGGTT	TAAAAGCGTGGCA			M.leprae
1098	CAGCCAAACTCCGAATGCCG-TGGTG-TATA	TCGTGGCA			M.gastri
1041	CAGCCAAACTCCGAATGCCG-TGGTG-TATA	TCGTGGCA			M.kansasii
3486	CAGCCAAACTCCGAATGCCG	ATAAGTAAAGATGCGA			M.smegmatis

Figure 1D

5/31




Figure 1E

6/31




Figure 1F

7/31

	1610	1620	1630	1640	
2558	GAT-CGATCAC-TCCCCTTCGGGGG	TGTGGAGTTC-TGG			<i>M. tuberculosis</i>
1860	GAT-CGACCAT-TCCCCTTCGGGGG	GTGGCGATT-CGG			<i>M. avium</i>
1860	GAT-CGACCAT-TCCCCTTCGGGGG	GTGGCGATT-CGG			<i>M. paratuberc.</i>
1955	GGC-CGATC-ATCC-TTCGGGG	GTGACGGTTG-GG			<i>M. phlei</i>
1879	GAT-CGACCAT-TCCCCTTCGGGGG	TATGGAGGTT-CGG			<i>M. leprae</i>
1641	GAT-CGATCAC-TCCCCTTCGGGGG	A-GTGGAGGTC-TGG			<i>M. gastri</i>
1584	GAT-CGATCAC-TCCCCTTCGGGGG	GTGGAGGTC-TGG			<i>M. kansasii</i>
4035	ACCGTGAACGCACT-TTCGGGG	TGTGGGTTGGTGG			<i>M. smegmatis</i>
	1650	1660	1670	1680	
2594	GGCTGCGTGGGAACCTTCGCTGGTAGTAGTCAAGC	GAAEGG			<i>M. tuberculosis</i>
1896	GGCTGCGTGGGAACCTTCGCTGGTAGTAGTCAAGC	AAATGGG			<i>M. avium</i>
1896	GGCTGCGTGGGAACCTTCGCTGGTAGTAGTCAAGC	AAATGGG			<i>M. paratuberc.</i>
1986	GGCTGCGTGGGAACCTTCGCTGGTAGTAGTCAAGC	GATGGG			<i>M. phlei</i>
1917	GGCTGCGTGGGAACCTTCGCTGGTAGTAGTCAAGC	GATGGG			<i>M. leprae</i>
1677	GGCTGCGTGGGAGCCTTCGCTGGTAGTAGTCAAGC	GATGGG			<i>M. gastri</i>
1620	GGCTGCGTGGGAGCCTTCGCTGGTAGTAGTCAAGC	GATGGG			<i>M. kansasii</i>
4071	GGCTGCAATGGGAACCTTCGCTGGTAGTAGTCAAGC	GATGGG			<i>M. smegmatis</i>
	1690	1700	1710	1720	
2634	-GTGACGCAGGAAGGTAGCCGTACCAGTCAGTGGTAATCA-				<i>M. tuberculosis</i>
1936	-GTGACGCAGGAAGGTAGCCGTACCAGTCAGTGGTAATA-				<i>M. avium</i>
1936	-GTGACGCAGGAAGGTAGCCGTACCAGTCAGTGGTAATA-				<i>M. paratuberc.</i>
2025	-GTGACGCAGGAAGGTAGCCGTACCAGTCAGTGGTAATA-				<i>M. phlei</i>
1957	-GTGACGCAGGAAGGTAGCCGTACCAGTCAGTGGTAATA-				<i>M. leprae</i>
1717	-GTGACGCAGGAAGGTAGCCGTACCAGTCAGTGGTAATA-				<i>M. gastri</i>
1660	-GTGACGCAGGAAGGTAGCCGTACCAGTCAGTGGTAATA-				<i>M. kansasii</i>
4111	-GTGACGCAGGAAGGTAGCCGTACCGTTCAGTGGTAATA-				<i>M. smegmatis</i>

Figure 1G

8/31

	1730	1740	1750	1760	
2672	-CTGGGGCAAGCC	GGTAGGG	AGAGCGATAGGCAAATCCGT		M.tuberculosis
1974	-CTGGGGCAAGCC	CGTAG	AGAGCGATAGGCAAATCCGT		M.avium
1974	-CTGGGGCAAGCC	CGTAG	AGAGCGATAGGCAAATCCGT		M.paratuberc.
2063	-C	GGGGTAA	CCGTAGGG	GGAGT	GATAGGCAAATCCGT M.phlei
1995	-CTGG	AGCAAGCC	GTAGGGAGAGCGATAGGCAAATCCGT		M.leprae
1755	-CTGGGGCAAGCC	AGTAGGGAGAGCGATAGGCAAATCCGT			M.gastri
1698	-CTGGGGCAAGCC	AGTAGGGAGAGCGATAGGCAAATCCGT			M.kansasii
4149	-C	GGG	GTAAAGCC	GTAGGGAGT	TCAGATAGGTAATCCGT M.smegmatis

- - - -

	1970	1980	1990	2000		
2908	AGGGGG	ACCGGAATAT	CGTGAACACCCTTGCGGTGGGAGC		M.tuberculosis	
2208	AGGGGG	CCGGAATA	CGTGAACACCCTTGCGGTGGGAGC		M.avium	
2208	AGGGGG	CCGGAATA	CGTGAACACCCTTGCGGTGGGAGC		M.paratuberc.	
2298	AGGGGG	ACCCACGTAC	CGTGA	GGGCTT	TGCGG	GGGAGC M.phlei
2231	AGGGGG	CCGGAATAT	CGTGAACACCCTTGCGGTGGGAGC		M.leprae	
1910					M.gastri	
1934	AGGGGG	ACCGGAATA	CGTGAACACCCTTGCGGTGGGAGC		M.kansasii	
4385	AGGGGG	ACCCACAT	GGCGTGT	TAAGCC	TTTACGG	CCCAAGC M.smegmatis

- - - -

	2410	2420	2430	2440	
3345	AC	CTCGACGCCAGTTGGGGC	GGAGTCGTTGTTGAAATACC		M.tuberculosis
284	ACCTCGACGCCAGTTGGGGC	GGAGTCGTTGTTGAAATACC			M.bovis
2645	GCACAGACGCCAGTTTGTG	TGGAGTCGTTGTTGAAATACC			M.avium
393	ATACAGACGCCAGTTTGTAT	TGGAGTCGTTGTTGAAATACC			M.intracellulare
2645	GCACAGACGCCAGTTTGTG	TGGAGTCGTTGTTGAAATACC			M.paratuberc.
2737	GCTCGGACGCCAGTTGGGT	TGGAGTCGTTGTTGAAATACC			M.phlei
2668	ACITCGACGC	TAGTTGGGGT	TGGAGTCGTTGTTGAAATACC		M.leprae
1910					M.gastri
2372	ACCTCA	ACGCCAGTTGGGGT	TGGAGTCGTTGTTGAAATACC		M.kansasii
4822	GCTCAC	ACGCCAGTTGGGT	TGGAGTCGTTGTTGAAATACC		M.smegmatis

Figure 1H

9/31

	2450	2460	2470	2480	
3385	ACTCTGATCGTATTGG	GCATCTAACCTCGAACCCCTGAATC			M.tuberculosis
324	ACTCTGATCGTATTGGG	CATCTAACCTCGAACCCCTGAATC			M.bovis
2685	ACTCTGATCGTATTGG	CACTAACCTCGAACCCCTTATC			M.avium
433	ACTCTGATCGTATTGG	CACTAACCTCGAACCCCTTATC			M.intracellulare
2685	ACTCTGATCGTATTGG	CACTAACCTCGAACCCCTTATC			M.paratuberc.
2777	ACTCTGATCGTATTGGG	CTAACCTCGAACCCCTGGATC			M.phlei
2708	ACTCTGAT	GTATTGACATCTAACCTCGAACCCCTATATC			M.leprae
1910					M.gastri
2412	ACTCTGATCGTATTGG	CACTAACCTCGAACCCCTGAATC			M.kansasii
4862	ACTCTGATCGTATTGGG	CTAACCTCGAACCCCTATATC			M.smegmatis

	2490	2500	2510	2520	
3425	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.tuberculosis
364	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.bovis
2724	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.avium
472	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.intracellulare
2724	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.paratuberc.
2817	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.phlei
2748	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.leprae
1910					M.gastri
2452	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.kansasii
4902	GGGTTTAG	GGACAGTGCCTGGCGGGTAGTTTAACTGGGGC			M.smegmatis

- - - -

	2930	2940	2950	2960	
3864	AGTACGAGAGGACCGGGACGGACGAACCTCT	GGTGACCA			M.tuberculosis
3163	AGTACGAGAGGACCGGGACGGACGAACCTCT	GGTATACCA			M.avium
3163	AGTACGAGAGGACCGGGACGGACGAACCTCT	GGTATACCA			M.paratuberc.
3256	AGTACGAGAGGACCGGGACGGACGAACCTCT	GGTATACCA			M.phlei
3187	AGTACGAGAGGACCGGGACGGACGAACCTCT	GGTATACCA			M.leprae
1910					M.gastri
2891	AGTACGAGAGGACCGGGACGGACGAACCTCT	GTGCACCA			M.kansasii
5342	AGTACGAGAGGACCGGGACGGACGAACCTCT	GGTATACCA			M.smegmatis

Figure 11

10/31

	2970	2980	2990	3000	
3904	GTTGT	CCCG	CCAGGGGCACCGCTGGATAGCCACGTTTCGGT		M.tuberculosis
3203	GTTGT	CCCG	CCAGGGGCACCGCTGGATAGCCACGTTTCGGT		M.avium
3203	GTTGT	CCCG	CCAGGGGCACCGCTGGATAGCCACGTTTCGGT		M.paratuberc.
3296	GTTGT	CCCG	CCAGGGGCACCGCTGGATAGCCACGTTTCGGT		M.phlei
3227	GTTGT	CTCA	CCAGGGGCACCGCTGGATAGCCACGTTTCGGT		M.leprae
1910					M.gastri
2931	GTTGT	CCCG	CCAGGGGCACCGCTGGATAGCTACGTTTCGGT		M.kansasii
5382	GTTGT	CCCG	CCAGGGGCACCGCTGGATAGCCACGTTTCGGT		M.smegmatis
	3010	3020	3030	3040	
3944	CAGGATAACCGCTGAAAGCATCTAAGCGGGAAACCTTCTC				M.tuberculosis
3243	CAGGATAACCGCTGAAAGCATCTAAGCGGGAAACCTTCTC				M.avium
3243	CAGGATAACCGCTGAAAGCATCTAAGCGGGAAACCTTCTC				M.paratuberc.
3336	CAGGATAACCGCTGAAAGCATCTAAGCGGGAAACCTCTTC				M.phlei
3267	CAGGATAACCGCTGAAAGCATCTAAGCGGGAAACCTTCTC				M.leprae
1910					M.gastri
2971	CAGGATAACCGCTGAAAGCATCTAAGCGGGAAACCTTCTC				M.kansasii
5422	CAGGATAACCGCTGAAAGCATCTAAGCGGGAAACCTCTTC				M.smegmatis
- - - -					
	3090	3100	3110	3120	
4023	CCCGC-AGAACACGGGTTCAATAGGTCAGACCTGGAAGCT				M.tuberculosis
609	CCCGC-AGAACACGGGTTCAATAGGTCAGACCTGGAAGCT				M.bovis
3322	CCCGC-AGAACACGGGTTGATAGGTCAGACCTGGAAGCT				M.avium
677	CCCGC-AGAACACGGGTTGATAGGTCAGACCTGGAAGCT				M.intracellulare
3322	CCCGC-AGAACACGGGTTGATAGGTCAGACCTGGAAGCT				M.paratuberc.
3415	CCCGC-AGAACACGGGTTGATAGGTCAGACCTGGAAGCT				M.phlei
3309					M.leprae
1910					M.gastri
3050	CCCGC-AGAACACGGGTTGATAGGTCAGACCTGGAAGCT				M.kansasii
5501	CCCGC-AGAACACGGGTTGATAGGTCAGACCTGGAAGCT				M.smegmatis

Figure 1J

11/31

		50	60	70	80	
2	GCGGCGTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.tuberculosis
141	GCGGCGTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.bovis
39	GCGGCGTACTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.avium
1	-----TTAACACATGCAAGTGAACGGAAAGGTCTC					M.intracellulare
39	GCGGCGTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.paratuberc.
2	GCGGCGTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.scrofulaceum
40	GCGGCGTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.leprae
2	GCGGCGTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.kansasii
2	GCGGCGTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.gastri
40	GCGGCGTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.gordonae
1	-----GTGCTTAACACATGCAAGTCGAACGGAAAGGTCTC					M.marinum
		90	100	110	120	
42	T-----TCGGAGATACTCGAGTGGCGAACGGGT					M.tuberculosis
181	T-----TCGGAGATACTCGAGTGGCGAACGGGT					M.bovis
79	T-----TCGGAGTACTCGAGTGGCGAACGGGT					M.avium
32	T-----TCGGAGTACTCGAGTGGCGAACGGGT					M.intracellulare
79	T-----TCGGAGTACTCGAGTGGCGAACGGGT					M.paratuberc.
42	T-----TCGGAGTACTCGAGTGGCGAACGGGT					M.scrofulaceum
80	TAAAAAATCTTTTATAGATACTCGAGTGGCGAACGGGT					M.leprae
41	T-----TCGGAGTACTCGAGTGGCGAACGGGT					M.kansasii
42	T-----TCGGAGTACTCGAGTGGCGAACGGGT					M.gastri
80	-----GGGTACTCGAGTGGCGAACGGGT					M.gordonae
36	T-----TCGGAGATACTCGAGTGGCGAACGGGT					M.marinum
		130	140	150	160	
70	GAGTAACACGTGGGTGATCTGCCCTGCACTTC-GGGATAA					M.tuberculosis
209	GAGTAACACGTGGGTGATCTGCCCTGCACTTC-GGGATAA					M.bovis
107	GAGTAACACGTGGGCAATCTGCCCTGCACTTC-GGGATAA					M.avium
59	GAGTAACACGTGGGCAATCTGCCCTGCACTTC-GGGATAA					M.intracellulare
107	GAGTAACACGTGGGCAATCTGCCCTGCACTTC-GGGATAA					M.paratuberc.
70	GAGTAACACGTGGGCAATCTGCCCTGCACTTC-GGGATAA					M.scrofulaceum
120	GAGTAACACGTGGGTATCTGCCCTGCACTTCGGGATAA					M.leprae
69	GAGTAACACGTGGGCAATCTGCCCTGCACACC-GGGATAA					M.kansasii
70	GAGTAACACGTGGGCAATCTGCCCTGCACACC-GGGATAA					M.gastri
104	GAGTAACACGTGGGTATCTGCCCTGCACATC-GGGATAA					M.gordonae
64	GAGTAACACGTGGGTGATCTGCCCTGCACTTC-GGGATAA					M.marinum

Figure 2A

12/31

	170	180	190	200	
109	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CACGGGA			M.tuberculosis
248	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CACGGGA			M.bovis
146	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CTCAAGA			M.avium
98	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CTTTAGG			M.intracellulare
146	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CTCAAGA			M.paratuberc.
109	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CACTTGG			M.scrofulaceum
160	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	TTCAAGG			M.leprae
108	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CACTTGG			M.kansasii
109	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CACTTGG			M.gastri
143	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CAAGGA			M.gordonae
103	GCCTGGGAAACTGGGTCTAATACCGGATAGGAC	CACGGGA			M.marinum
	210	220	230	240	
149	TGCATGTCTTGTGGTGGAAAG	CGCTTTAG	CGGTGTGGGAT		M.tuberculosis
288	TGCATGTCTTGTGGTGGAAAG	CGCTTTAG	CGGTGTGGGAT		M.bovis
186	CGCATGTCTTGTGGTGGAAAG	CTTTT	ACGGTGTGGGAT		M.avium
138	CGCATGTCTTGTGGTGGAAAG	CTTTT	GCGGTGTGGGAT		M.intracellulare
186	CGCATGTCTTGTGGTGGAAAG	CTTTT	GCGGTGTGGGAT		M.paratuberc.
149	CGCATGTCTTGTGGTGGAAAG	CTTTT	GCGGTGTGGGAT		M.scrofulaceum
200	CGCATGTCTTGTGGTGGAAAG	CTTTT	GCGGTGTGGGAT		M.leprae
148	CGCATGTCTTGTGGTGGAAAG	CTTTT	GCGGTGTGGGAT		M.kansasii
149	CGCATGTCTTGTGGTGGAAAG	CTTTT	GCGGTGTGGGAT		M.gastri
183	CACATGTCTTGTGGTGGAAAG	CTTTT	GCGGTGTGGGAT		M.gordonae
143	TTCATGTCTTGTGGTGGAAAG	CTTTT	GCGGTGTGGGAT		M.marinum
	250	260	270	280	
189	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.tuberculosis
328	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.bovis
224	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.avium
176	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.intracellulare
224	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.paratuberc.
187	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.scrofulaceum
239	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.leprae
186	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.kansasii
187	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.gastri
221	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.gordonae
181	GAGCCCGCGGCCTATCAGCTTGTGGTGGGGTG	ACGGCCT			M.marinum

Figure 2B

13/31

450 460 470 480

389 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.tuberculosis
 528 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.bovis
 424 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.avium
 376 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.intracellulare
 424 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.paratuberc.
 387 AAACCTCTTTCACCATCGACGAAGGTCGA---CTTGTGG M.scrofulaceum
 439 AAACCTCTTTCACCATCGACGAAGGTCGGGAATCTCTCGG M.leprae
 386 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.kansasii
 387 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.gastri
 420 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.gordonae
 381 AAACCTCTTTCACCATCGACGAAGGTCGGGGTTCTCTCGG M.marinum

- - - -

1130 1140 1150 1160

1069 TCTCATGTTGCCAGCACGTAATGGTGGGGACTCGTGAGAG M.tuberculosis
 1208 TCTCATGTTGCCAGCACGTAATGGTGGGGACTCGTGAGAG M.bovis
 1104 TCTCATGTTGCCAGCGGGTAATGCGGGGACTCGTGAGAG M.avium
 1056 TCTCATGTTGCCAGCGGGTAATGCGGGGACTCGTGAGAG M.intracellulare
 1098 TCTCATGTTGCCAGCGGGTAATGCGGGGACTCGTGAGAG M.paratuberc.
 1064 TCTCATGTTGCCAGCGGGTAATGCGGGGACTCGTGAGAG M.scrofulaceum
 1119 TCTCATGTTGCCAGCACGTAATGGTGGGGACTCGTGAGAG M.leprae
 1066 TCTCATGTTGCCAGCGGGTAATGCGGGGACTCGTGAGAG M.kansasii
 1067 TCTCATGTTGCCAGCGGGTAATGCGGGGACTCGTGAGAG M.gastri
 1100 TCTCATGTTGCCAGCGGGTAATGCGGGGACTCGTGAGAG M.gordonae
 1061 TCTCATGTTGCCAGCACGTAATGGTGGGGACTCGTGAGAG M.marinum

- - - -

1250 1260 1270 1280

1189 CAATGGCCGGTACAAAGGGCTGCGATGCCGCGAGGTTAAG M.tuberculosis
 1328 CAATGGCCGGTACAAAGGGCTGCGATGCCGCGAGGTTAAG M.bovis
 1224 CAATGGCCGGTACAAAGGGCTGCGATGCCGTAAGGTTAAG M.avium
 1176 CAATGGCCGGTACAAAGGGCTGCGATGCCGTAAGGTTAAG M.intracellulare
 1218 CAATGGCCGGTACAAAGGGCTGCGATGCCGTAAGGTTAAG M.paratuberc.
 1184 CAATGGCCGGTACAAAGGGCTGCGATGCCGTAAGGTTAAG M.scrofulaceum
 1239 CAATGGCCGGTACAAAGGGCTGCGATGCCGTAAGGTTAAG M.leprae
 1186 CAATGGCCGGTACAAAGGGCTGCGATGCCGCGAGGTTAAG M.kansasii
 1187 CAATGGCCGGTACAAAGGGCTGCGATGCCGCGAGGTTAAG M.gastri
 1220 CAATGGCCGGTACAAAGGGCTGCGATGCCGCGAGGTTAAG M.gordonae
 1181 CAATGGCCGGTACAAAGGGCTGCGATGCCGCGAGGTTAAG M.marinum

Figure 2C

14/31

	1290	1300	1310	1320	
1229	CGAATCCTTA-AAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.tuberculosis
1368	CGAATCCTTA-AAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.bovis
1264	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.avium
1216	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.intracellulare
1258	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.paratuberc.
1224	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.scrofulaceum
1279	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.leprae
1226	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.kansasii
1227	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.gastri
1260	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.gordonae
1221	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGATCGGGGTCT				M.marinum

	1330	1340	1350	1360	
1268	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.tuberculosis
1407	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.bovis
1304	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.avium
1256	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.intracellulare
1298	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.paratuberc.
1264	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.scrofulaceum
1319	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.leprae
1266	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.kansasii
1267	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.gastri
1300	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.gordonae
1260	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.marinum

Figure 2D

15/31

		50	60	70	80	
128	TTCCGAACCCGGAAGCTAAGCCTGCCAGCGCCGATGATAC					M.tuberculosis
39	TGCCGAACCCGGAAGCTAAGCCTGCCAGCGCCGATGATAC					M.bovis
41	TGCCGAACCCGGAAGCTAAGCCTGCCAGCGCCGATGATAC					M.phlei
3559	TACCGAACCCGGAAGCTAAGCCTGCCAGCGCCGATGATAC					M.leprae
5743	TGCCGAACCCGGAAGCTAAGCCTGCCAGCGCCGATGATAC					M.smegmatis
		90	100	110	120	
168	TGCCCCTCCGGG---TGGAAAAGTAGGACACCGCCGAAC					M.tuberculosis
79	TGCCCCTCCGGG---TGGAAAAGTAGGACACCGCCGAAC					M.bovis
81	TGCCCCTCACCGGG---TGGAAAAGTAGGACACCGCCGAAC					M.phlei
3599	TGCCCCTTCGGG---TGGAAAAGTAGGACACCGCCGAAC					M.leprae
5782	TACCCCTTCGGG---TGGAAAAGTAGGACACCGCCGAAC					M.smegmatis

Figure 3

16/31

	90	100	110	120	
382	GGGAGCTGTCAACCGAGC	ATTGATCCGAGGATTTCCGAAT			M.avium
382	GGGAGCTGTCAACCGAGC	ATTGATCCGAGGATTTCCGAAT			M.paratuberc.
1053	GGGAGCTGTCAACCGAGC	TTGGATCCGAGGATTTCCGAAT			M.tuberculosis
467	GGGAGCTGTCAACCGAGC	TTGGATCCGAGGATTTCCGAAT			M.phlei
392	GGGAGCTGTCAACCGAGC	TTGGATCCGAGGATTTCCGAAT			M.leprae
167	GGGAGCTGTCAACCGAGC	TTGGATCCGAGGATTTCCGAAT			M.gastri
110	GGGAGCTGTCAACCGAGC	TTGGATCCGAGGATTTCCGAAT			M.kansasii
2548	GGGAGCTGTCAACCGAGC	TTGATCCGAGGATTTCCGAAT			M.smegmatis
- - - -					
	170	180	190	200	
462	GAATATATAGGGTGCG-GGAGG	TAAACGCGGGGAAGTGAAA			M.avium
462	GAATATATAGGGTGCG-GGAGG	TAAACGCGGGGAAGTGAAA			M.paratuberc.
1133	GAATATATAGGGTGCG-GGAGG	TAAACGCGGGGAAGTGAAA			M.tuberculosis
547	GAATATATAGGCGTTG-GGG	GGGAACGCGGGGAAGTGAAA			M.phlei
472	GAATATATAGGGTGCG-GGAGG	TAAACGCGGGGAAGTGAAA			M.leprae
247	GAATATATAGGGTGCG-GGAGG	TAAACGCGGGGAAGTGAAA			M.gastri
190	GAATATATAGGGTGCG-GGAGG	TAAACGCGGGGAAGTGAAA			M.kansasii
2628	GAATATATAGGCGTCT-GGG	GGGAACGCGGGAAGTGAAA			M.smegmatis
- - - -					
	250	260	270	280	
541	-GTCAGTAGTGGCGAGCGAAC	-CGGAACA-GGCTAAACCG			M.avium
541	-GTCAGTAGTGGCGAGCGAAC	-CGGAACA-GGCTAAACCG			M.paratuberc.
1212	-GTCAGTAGTGGCGAGCGAAC	-CGGAACA-GGCTAAACCG			M.tuberculosis
626	-GTCAGTAGTGGCGAGCGAAC	-AGGGAGGATGGCTAAACCG			M.phlei
551	-GTCAGTAGTGGCGAGCGAAC	-GTGGAAATATGGCTAAACCG			M.leprae
326	-GTCAGTAGTGGCGAGCGAAC	-CGGAACATGGCTAAACCG			M.gastri
269	-GTCAGTAGTGGCGAGCGAAC	-CGGAACATGGCTAAACCG			M.kansasii
2706	-GTCAGTAGTGGCGAGCGAAC	-CGGAGGATGGCTAAACG			M.smegmatis

Figure 4A

17/31

	290	300	310	320	
578	CATG	CATG	GACA	ACCGGGTAGGGGTTGTGTGTGCGGGGT	M. avium
578	CATG	CATG	GACA	ACCGGGTAGGGGTTGTGTGTGCGGGGT	M. paratuberc.
1250	CACG	CATG	GTA	ACCGGGTAGGGGTTGTGTGTGCGGGGT	M. tuberculosis
664	CGTG	CATG	TGATA	CCCGGTGGGGTTGTGTGTGCGGGGT	M. phlei
590	CACA	CATG	TCTA	ACTAGGTAGGGGTTGTGTGTGCGGGGT	M. leprae
365	CACG	CATG	GGTG	ACCGGGTAGGGGTTGTGTGTGCGGGGT	M. gastri
308	CACG	CATG	GGTG	ACCGGGTAGGGGTTGTGTGTGCGGGGT	M. kansasii
2745	TATGA	CATG	TGATA	ACCGGGTAGGGGTTGTGTGTGCGGGGT	M. smegmatis

	330	340	350	360	
617	TGTGGG	ATTGATATG	TCTCAGCT	TCTACCTGGCTGAGG	-GG M. avium
617	TGTGGG	ATTGATATG	TCTCAGCT	TCTACCTGGCTGAGG	-GG M. paratuberc.
1289	TGTGGG	GATATGTCTCAGCG	CTACCGGCTGAGG	-GG M. tuberculosis	
703	TGTGGG	CCCTGTGTGTC	CATCGTCCGCGGGC	GATGGCAG M. phlei	
629	TGTGGG	ATTGTATGTCTCA	CTCTACCTGGTTGAGG	-GG M. leprae	
404	TGTGGG	ATTGATA	GTCTCAGCTCTACCGGCTGAGG	-GG M. gastri	
347	TGTGGG	ATTGATA	GTCTCAGCTCTACCGGCTGAGG	-GG M. kansasii	
2785	TGTGGG	ACCTATCTTTC	CGCTCTACCTGGCTG	GAGG M. smegmatis	

	370	380	390	400	
656	TAGTCAGAAA	GTGTCGTGGTTAGCGGAAGTGGCCTGGG	ATC	M. avium	
656	TAGTCAGAAA	GTGTCGTGGTTAGCGGAAGTGGCCTGGG	AC	M. paratuberc.	
1327	CAGTCAGAAA	GTGTCGTGGTTAGCGGAAGTGGCCTGGG	AT	M. tuberculosis	
742	TAGTGA	ATAAAGCAGTGTGGTTAGGTGAAGTGGCCTGGG	AT	M. phlei	
668	TAGTCAGAAA	GTGTCGTGGTTAGCGGAAGTGGCCTGGG	AT	M. leprae	
443	CAGTCAGAAA	GTGTCGTGGTTAGCGGAAGTGGCCTGGG	AT	M. gastri	
386	CAGTCAGAAA	GTGTCGTGGTTAGCGGAAGTGGCCTGGG	AT	M. kansasii	
2823	CAGTGA	AGAAAATGTTGTGGTTAGCGGAAGTGGC	ATGGGAT	M. smegmatis	

Figure 4B

18/31

	410	420	430	440	
696	GGCCCC	GCCGTAGACGGT	GAGAGCCCCGGT	TACGCGAAA-ACC	M.avium
696	GGCCCC	GCCGTAGACGGT	GAGAGCCCCGGT	TACGCGAAA-ACC	M.paratuberc.
1367	GGTCTG	GCCGTAGACGGT	GAGAGCCCCGGT	TACGCGAAA-ACC	M.tuberculosis
782	GGTCTG	GCCGTAGTGGT	GAGAGCCCCGTAAC	CGAAA-ACA	M.phlei
708	GGCCTG	GCCGTAGACGGT	GAGAGCCCCGTACGCG	AAA-ACC	M.leprae
483	GGTCTG	GCCGTAGACGGT	GAGAGCCCCGGTACGT	GAAA-ACC	M.gastri
426	GGTCTG	GCCGTAGACGGT	GAGAGCCCCGGTACGT	GAAA-ACC	M.kansasii
2863	GGCCTG	GCCGTAGACGGT	GAGAGCCCCGGTACGT	GAAA-ACC	M.smegmatis

	450	460	470	480	
735	CGGCACCTGCCCT	TATATCAACA	CCGAGTAGCAGCGGGCC		M.avium
735	CGGCACCTGCCCT	TATATCAACACCCG	AGTAGCAGCGGGCC		M.paratuberc.
1406	CGGCACCTGCCCT	AGTATCAATT	CCCGAGTAGCAGCGGGCC		M.tuberculosis
820	TGCTGCC	GCCTGCACAGG--	TCCCGAGTAGCAGCGGGCC		M.phlei
747	TGGCACCTGCCCT	TGATCAATT	CCCGAGTAGCAGCGGGCC		M.leprae
522	CGGCACCTGCCCT	TGATCAATT	CCCGAGTAGCAGCGGGCC		M.gastri
465	CGGCACCTGCCCT	TGATCAATT	CCCGAGTAGCAGCGGGCC		M.kansasii
2902	CGACGTCTG	CTTGATGGTGT	CCCGAGTAGCAGCGGGCC		M.smegmatis

- - - -

	570	580	590	600	
855	GAGGGAATGGT	GAAAAGTACCCCGGG	AGGG-AGTGAAATA		M.avium
855	GAGGGAATGGT	GAAAAGTACCCCGGGAGGG	-AGTGAAATA		M.paratuberc.
1526	GAGGGAATGGT	GAAAAGTACCCCGGGAGGG	AGTGAAAGA		M.tuberculosis
937	GAGGGAAT	GTGAAAAGTACCCCGGGAGGG	-AGTGAAAGA		M.phlei
867	GAGGGAATGGT	GAAAAGTACCCCGGGAGGG	AGTGAAATA		M.leprae
642	GAGGGAATGGT	GAAAAGTACCCCGGGAGGG	AGTGAAAGA		M.gastri
585	GAGGGAATGGT	GAAAAGTACCCCGGGAGGG	AGTGAAAGA		M.kansasii
3022	GAGGGAATGGT	GAAAAGTACCCCGGGAGGG	AGTGAAAGA		M.smegmatis

Figure 4C

19/31

	610	620	630	640	
894	GTACCTGAAACCGTGTGCCTACAATCCGTCAGAGCC	TCCT			M.avium
894	GTACCTGAAACCGTGTGCCTACAATCCGTCAGAGCCTCCT				M.paratuberc.
1566	GTACCTGAAACCGTGTGCCTACAATCCGTCAGAGCCTCCT				M.tuberculosis
976	GTACCTGAAACCGTGTGCCTACAATCCGTCAGAGCC	CTCT			M.phlei
907	GTACCTGAAACCGTGTGCCTACAATCCGTCAGAGCCTCCT				M.leprae
682	GTACCTGAAACCGTGTGCCTACAATCCGTCAGAGCC	CTCT			M.gastri
625	GTACCTGAAACCGTGTGCCTACAATCCGTCAGAGCC	CTTT			M.kansasii
3062	GTACCTGAAACCGTGTGCCTACAATCCGTCAGAGCC	CTCG			M.smegmatis

	650	660	670	680	
934	C-----GTGGGGTGATGGCGTGCCTTTTGA				M.avium
934	C-----GTGGGGTGATGGCGTGCCTTTTGA				M.paratuberc.
1606	TTTCCTCTCCGGAGGAGGGTGGTGATGGCGTGCCTTTTGA				M.tuberculosis
1016	CTT-----GTAGTGGGGTGATGGCGTGCCTTTTGA				M.phlei
947	T-----GTGGGGTGATGGCGTGCCTTTTGA				M.leprae
722	T-----GTGGGGTGATGGCGTGCCTTTTGA				M.gastri
665	C-----GTGGGGTGATGGCGTGCCTTTTGA				M.kansasii
3102	ACGTGT-----GTGGGGTGATGGCGTGCCTTTTGA				M.smegmatis

	690	700	710	720	
959	AGAATGAGCCTGCGAGTCAGGGT	CACGTCGCG	AGGTTAAC		M.avium
23	AGAATGAGCCTGCGAGTCAGGGACACGTCGCGAGGTTAAC				M.intracellulare
959	AGAATGAGCCTGCGAGTCAGGGACACGTCGCGAGGTTAAC				M.paratuberc.
1646	AGAATGAGCCTGCGAGTCAGGGACATGTCGCGAGGTTAAC				M.tuberculosis
4	AGAATGAGCCTGCGAGTCAGGGACATGTCGCGAGGTTAAC				M.bovis
1046	AGAATGAGCCTGCGAGTCAGGGACATGTCGCGAGGTTAAC				M.phlei
972	AGAATGAGCCTGCGAGTCAGGGACATGTCGCGAGGTTAAC				M.leprae
747	AGAATGAGCCTGCGAGTCAGGGACATGTCGCGAGGTTAAC				M.gastri
690	AGAATGAGCCTGCGAGTCAGGGACATGTCGCGAGGTTAAC				M.kansasii
3132	AGAATGAGCCTGCGAGTCAGGGACATGTCGCGAGGTTAAC				M.smegmatis

Figure 4D

20/31

770 780 790 800

1039 CGCATCCCCTTTGGG-----GTGTAGTGGCGTGT M.avium
 103 CGCATCCCCTTTGGG-----GTGTAGTGGCGTGT M.intracellulare
 1039 CGCATCCCCTTTGGG-----GTGTAGTGGCGTGT M.paratuberc.
 1726 CGACCCACACGCGCATACGCGCGTGTGAATAGTGGCGTGT M.tuberculosis
 84 CGACCCACACGCGCATACGCGCGTGTGAATAGTGGCGTGT M.bovis
 1126 CGTATCCAACTGTGTT----GGGGTTGGTGTAGTGGTGTGT M.phlei
 1052 CGTATCACGTGTGAGCGT-----GTGTAGTGGCGTGT M.leprae
 827 CGTATCACGCGTAAGCGT-----GTGTAGTGGCGTGT M.gastri
 770 CGTATCGCGCGGAGCGT-----GTGTAGTGGCGTGT M.kansasii
 3212 CGTATCCACACAAGAGTGTGTG----GTGTAGTGGTGTGT M.smegmatis

- - - -

1050 1060 1070 1080

1307 CAGCCAAACTCCGAATGCCG-TGGTG-TAAAAGCGTGGCA M.avium
 1307 CAGCCAAACTCCGAATGCCG-TGGTG-TAAAAGCGTGGCA M.paratuberc.
 2005 CAGCCAAACTCCGAATGCCG-TGGTG-TA AAGCGTGGCA M.tuberculosis
 1401 CAGCCAAACTCCGAATGCCGTAAG-TAAAAGTGTGGCA M.phlei
 1323 CAGCCAAACTCCGAATGCCG-TGGTT-TAAAAGCGTGGCA M.leprae
 1098 CAGCCAAACTCCGAATGCCG-TGGTG-TATA GCGTGGCA M.gastri
 1041 CAGCCAAACTCCGAATGCCG-TGGTG-TATA GCGTGGCA M.kansasii
 3486 CAGCCAAACTCCGAATGCCGTAAGGCCAAGAGTGGGTA M.smegmatis

- - - -

1170 1180 1190 1200

1425 AGTGGAAAAGGATGTGTAGTCGCAGAGACAAACCAGGAGG M.avium
 1425 AGTGGAAAAGGATGTGTAGTCGCAGAGACAAACCAGGAGG M.paratuberc.
 2122 AGTGGGAAAGGATGTGCAGTCGCAGAGACAAACCAGGAGG M.tuberculosis
 1519 AGTGGAAAAGGATGTGCAGTCGCAGAGACAAACCAGGAGG M.phlei
 1441 AGTGGGAAAGGATGTGCAGTCGCAGAGACAAACCAGGAGG M.leprae
 1215 AGTGGGAAAGGATGTGCAGTCGCAGAGACAAACCAGGAGG M.gastri
 1158 AGTGGGAAAGGATGTGCAGTCGCAGAGACAAACCAGGAGG M.kansasii
 3606 AGTGGAAAAGGATGTGTAGTCGCAGAGAGAAACCAGGAGG M.smegmatis

Figure 4E

21/31

	1250	1260	1270	1280	
1504	CTCACTGGTCAAGTGATT	ATGCGCC	ATAATGTAGCGGGG		M.avium
1504	CTCACTGGTCAAGTGATT	ATGCGCCGATAATGTAGCGGGG			M.paratuberc.
2201	CTCACTGGTCAAGTGATT	GTGCGCCGATAATGTAGCGGGG			M.tuberculosis
1598	CTCACTGGTCAAGTGATT	GTGCGC	ATAATGTAGCGGGG		M.phlei
1520	CTCACTGGTCAAGTGATT	GTGCGCCGATAATGTAGCGGGG			M.leprae
1294	CTCACTGGTCAAGTGATT	GTGCGCCGATAATGTAGCGGGG			M.gastri
1237	CTCACTGGTCAAGTGATT	GTGCGCCGATAATGTAGCGGGG			M.kansasii
3686	CTCACTGGTCAAGTGATT	GTGCGCCGATA	TGTGCGGGG		M.smegmatis

	1290	1300	1310	1320	
1544	CTCAAGCACACCGCCGAAGCCGCGGCACAT	TCATCTT-TA			M.avium
1544	CTCAAGCACACCGCCGAAGCCGCGGCACAT	TCATCTT-TA			M.paratuberc.
2241	CTCAAGCACACCGCCGAAGCCGCGGCACAT	CCACCTTGT			M.tuberculosis
1638	CTCAAGCACACCGCCGAAGCCGCGGCA	ATCAGCCTTTG			M.phlei
1560	CTCAAGCACACCGCCGAAGCCGCGGCACAT	TCACCTTCTA			M.leprae
1334	CTCAAGCACACCGCCGAAGCCGCGGACA	ACCGC	A		M.gastri
1277	CTCAAGCACACCGCCGAAGCCGCGGACA	ACCGC	A		M.kansasii
3726	CTCAAGCACACCGCCGAAGCCGCGGAA	GCCACCTTTG			M.smegmatis

	1330	1340	1350	1360	
1583	CGGTGGA	TGTGGGTAGGGGAGCGT	CCCC	CATTCAGCGAAG	M.avium
1583	CGGTGGATGTGGGTAGGGGAGCGT	CCCCCATTCAGCGAAG			M.paratuberc.
2280	GGTGGGTGTGGGTAGGGGAGCGT	CCCTCATTCAGCGAAG			M.tuberculosis
1676	TGGCTGGGTGTGGGTAGGGGAGCGT	CCGATCCGGGAAG			M.phlei
1600	GGGTGGATGTGGGTAGGGGAGCGT	CCCTCATTCAGCGAAG			M.leprae
1367	AGGT	TGGGTAGGGGAGCGT	CCCTCATTCAGCGAAG		M.gastri
1310	AGGT	TGGGTAGGGGAGCGT	CCCTCATTCAGCGAAG		M.kansasii
3764	TT	TGGGTAGGGGAGCGT	CCCTCATTCAGCGAAG		M.smegmatis

Figure 4F

22/31

1370 1380 1390 1400

1623 CT-CCGGGTGACCGGTGGTGGAGGGTGGGGGAGTGAGAAT M.avium
 1623 CT-CCGGGTGATCCGGTGGTGGAGGGTGGGGGAGTGAGAAT M.paratuberc.
 2319 CCACCGGGTGACCGGTGGTGGAGGGTGGGGGAGTGAGAAT M.tuberculosis
 1716 CCGCCGAGTGAACGGTGGTGGAGGGTGAGGAGTGAGAAT M.phlei
 1640 CTTCCGGGTACCGGTGGTGGAGGGTGGGGAGTGAGAAT M.leprae
 1402 CCGCCGGGTGACCGGTGGTGGAGGGTGGGGGAGTGAGAAT M.gastri
 1345 CTGCCGGGTGACCGGTGGTGGAGGGTGGGGGAGTGAGAAT M.kansasii
 3796 CCGCCGAGTATCGAGTGGTGGAGGGTGAGGAGTGAGAAT M.smegmatis

- - - -

1530 1540 1550 1560

1781 CGATGGACAACGGGTTGATATTCCTGATCCCGTGTATGGG M.avium
 1781 CGATGGACAACGGGTTGATATTCCTGATCCCGTGTATGGG M.paratuberc.
 2479 CGATGGACAACGGGTTGATATTCCTGATCCCGTGTGTGGG M.tuberculosis
 1875 CGATGGACAACGGGTTGATATTCCTGATCCCGTGTATGAG M.phlei
 1800 CGATGGACAACGGGTTGATATTCCTGATCCCGTGTGTGGG M.leprae
 1562 CGATGGACAACGGGTTGATATTCCTGATCCCGTGTGTGGG M.gastri
 1505 CGATGGACAACGGGTTGATATTCCTGATCCCGTGTGTGGG M.kansasii
 3956 CGATGGACAACGGGTTGATATTCCTGATCCCGTGTATGGG M.smegmatis

1570 1580 1590 1600

1821 CGTCCCTGATGAATCA-GCGGTACTAACCACCCAAAACCG M.avium
 1821 CGTCCCTGATGAATCA-GCGGTACTAACCACCCAAAACCG M.paratuberc.
 2519 CGTCCCTGATGAATCA-GCGGTACTAACCACCCAAAACCG M.tuberculosis
 1915 CGTCCCTGATGAATCA-GCGGTACTAACCACCCAAAACCG M.phlei
 1840 CGTCCCTGATGAATCA-GCGGTACTAACCACCCAAAACCG M.leprae
 1602 CGTCCCTGATGAATCA-GCGGTACTAACCACCCAAAACCG M.gastri
 1545 CGTCCCTGATGAATCA-GCGGTACTAACCACCCAAAACCG M.kansasii
 3996 CGTCCCTGATGAATCA-GCGGTACTAACCACCCAAAACCG M.smegmatis

Figure 4G

23/31

	1610	1620	1630	1640	
1860	GAT-CGATCCAT-TCCCTTCGGGGG	C-GTGGCGATT-CGG			M.avium
1860	GAT-CGACCAT-TCCCTTCGGGGG	C-GTGGCGATT-CGG			M.paratuberc.
2558	GAT-CGATCAQ-TCCCTTCGGGGG	TGTGGAGTTC-TGG			M.tuberculosis
1955	GSC-CGATC-ATCC-TTCGGGG	-GTGACGGTTG-GG			M.phlei
1879	GAT-CGACCATA-TCCCTTCGGGGG	CATGGAGGTT-CGG			M.leprae
1641	GAT-CGATCAC-TCCCTTCGGGGG	A-GTGGAGGTC-TGG			M.gastri
1584	GAT-CGATCAC-TCCCTTCGGGGG	C-GTGGAGGTC-TGG			M.kansasii
4035	ACCGTGACCGCACT-TTCGGGG	-TGTGGCGTTGGTGG			M.smegmatis
	1650	1660	1670	1680	
1896	GGCTGCGTGGGACCTTCGCT	TGGTAGTAGTCAAGCAATGGG			M.avium
1896	GGCTGCGTGGGACCTTCGCT	TGGTAGTAGTCAAGCAATGGG			M.paratuberc.
2594	GGCTGCGTGGGAACTTCGCT	TGGTAGTAGTCAAGCGATGGG			M.tuberculosis
1986	GGCTGCGTGGGACCG-GTGGG	TAGTAGTCAAGCGATGGG			M.phlei
1917	GGCTGCGTGGGAACTTCGCT	TGGTAGTAGTCAAGCGATGGG			M.leprae
1677	GGCTGCGTGGGAGCCTTCGCT	TGGTAGTAGTCAAGCGATGGG			M.gastri
1620	GGCTGCGTGGGAGCCTTCGCT	TGGTAGTAGTCAAGCGATGGG			M.kansasii
4071	GGCTGCGTGGGACCTTCGCT	TGGTAGTAGTCAAGCGATGGG			M.smegmatis
	1690	1700	1710	1720	
1936	-GTGACGCAGGAAGGCAGCCGT	TACCAGTCAGTGGTAATA-			M.avium
1936	-GTGACGCAGGAAGGCAGCCGT	TACCAGTCAGTGGTAATA-			M.paratuberc.
2634	-GTGACGCAGGAAGGTTAGCCGT	TACCAGTCAGTGGTAATA-			M.tuberculosis
2025	-GTGACGCAGGAAGGTTAGCCGT	TACCAGTCAGTGGTAATA-			M.phlei
1957	-GTGACGCAGGAAGGTTAGCCGT	TACCAGTCAGTGGTAATA-			M.leprae
1717	-GTGACGCAGGAAGGCAGCCGT	TACCAGTCAGTGGTAATA-			M.gastri
1660	-GTGACGCAGGAAGGCAGCCGT	TACCAGTCAGTGGTAATA-			M.kansasii
4111	-GTGACGCAGGAAGGTTAGCCGT	TACCAGTCAGTGGTAATA-			M.smegmatis
	1730	1740	1750	1760	
1974	-CTGGGGCAAGCCCGTAG--A	SAGCGATAGGCAAATCCGT			M.avium
1974	-CTGGGGCAAGCCCGTAG--A	SAGCGATAGGCAAATCCGT			M.paratuberc.
2672	-CTGGGGCAAGCCCGTAGGG	GAGAGCGATAGGCAAATCCGT			M.tuberculosis
2063	-CTGGGGCAAGCCCGTAGGG	GAGAGCGATAGGCAAATCCGT			M.phlei
1995	-CTGGGGCAAGCCCGTAGGG	GAGAGCGATAGGCAAATCCGT			M.leprae
1755	-CTGGGGCAAGCCCGTAGGG	GAGAGCGATAGGCAAATCCGT			M.gastri
1698	-CTGGGGCAAGCCCGTAGGG	GAGAGCGATAGGCAAATCCGT			M.kansasii
4149	-CTGGGGCAAGCCCGTAGGG	GAGAGCGATAGGCAAATCCGT			M.smegmatis

Figure 4H

24/31




Figure 4I

25/31

	2130	2140	2150	2160	
2367	CCGTTAACCC	GT	--AAGGGTGAAGCGGAGAATTTAAGCCC		M.avium
2367	CCGTTAACCC	GT	--AAGGGTGAAGCGGAGAATTTAAGCCC		M.paratuberc.
3067	CCGTTAACCC	GT	--AAGGGTGAAGCGGAGAATTTAAGCCC		M.tuberculosis
2457	CCGTTAACCC	TTTCGG	GGGTGAAGCGGAGAATTTAAGCCC		M.phlei
2390	CTGTTAACCC	GA	--AAGGGTGAAGCGGAGAATTTAAGCCC		M.leprae
1910					M.gastri
2094	CCGTTAACCC	GT	--AAGGGTGAAGCGGAGAATTTAAGCCC		M.kansasii
4544	CCGTTAACCC	CCTTGG	GGGTGAAGCGGAGAATTTAAGCCC		M.smegmatis

- - - -

	2250	2260	2270	2280	
2485	GTAACGACTTC	CCAA	CTGTCTCAACCATAGACTCGGCGAA		M.avium
2485	GTAACGACTTC	CCAA	CTGTCTCAACCATAGACTCGGCGAA		M.paratuberc.
3185	GTAACGACTTC	CAACTGTCTCAACCATAGACTCGGCGAA			M.tuberculosis
2577	GTAACGACTTC	CAACTGTCTCAACCATAGACTCGGCGAA			M.phlei
2508	GTAACGACTTC	CAACTGTCTCAACCATAGACTCGGCGAA			M.leprae
1910					M.gastri
2212	GTAACGACTTC	CAACTGTCTCAACCATAGACTCGGCGAA			M.kansasii
4663	GTAACGACTTC	CAACTGTCTCAAC	ATAGACTCGGCGAA		M.smegmatis

- - - -

	2370	2380	2390	2400	
2605	GTTTCGGTACGGTTTGTGTAGGATAGGTGGGAGACT	TTTGAA			M.avium
2605	GTTTCGGTACGGTTTGTGTAGGATAGGTGGGAGACT	TTTGAA			M.paratuberc.
3305	GTTTCGGTACGGTTTGTGTAGGATAGGTGGGAGACT	TTTGAA			M.tuberculosis
2697	GTTTCGGTACGGTTTGTGTAGGATAGGTGGGAGACT	TTTGAA			M.phlei
2628	GTTTCGGTACGGTTTGTGTAGGATAGGTGGGAGACT	TTTGAA			M.leprae
1910					M.gastri
2332	GTTTCGGTACGGTTTGTGTAGGATAGGTGGGAGACT	TTTGAA			M.kansasii
4782	GTTTCGGTACGGTTTGTGTAGGATAGGTGGGAGACT	TTTGAA			M.smegmatis

Figure 4J

26/31

	2410	2420	2430	2440	
2645	GCACAGACGCCAGTTTGGTGGAGTCGTTGTTGAAATACC				<i>M. avium</i>
393	ATACAGACGCCAGTTTGGTGGAGTCGTTGTTGAAATACC				<i>M. intracellulare</i>
2645	GCACAGACGCCAGTTTGGTGGAGTCGTTGTTGAAATACC				<i>M. paratuberc.</i>
3345	ACCTCGACGCCAGTTGGGGGGGAGTCGTTGTTGAAATACC				<i>M. tuberculosis</i>
284	ACCTCGACGCCAGTTGGGGGGGAGTCGTTGTTGAAATACC				<i>M. bovis</i>
2737	GCTCGACGCCAGTTGGGGGGGAGTCGTTGTTGAAATACC				<i>M. phlei</i>
2668	ACTTCGACGCTAGTTGGGGGGGAGTCGTTGTTGAAATACC				<i>M. leprae</i>
1910					<i>M. gastri</i>
2372	ACCTCAACGCCAGTTGGGGGGGAGTCGTTGTTGAAATACC				<i>M. kansasii</i>
4822	GCTCAACGCCAGTTGGGGGGGAGTCGTTGTTGAAATACC				<i>M. smegmatis</i>
	2450	2460	2470	2480	
2685	ACTCTGATCGTATTGGACACCTAACGTCGAACCCCT-TATC				<i>M. avium</i>
433	ACTCTGATCGTATTGGACACCTAACGTCGAACCCCT-TATC				<i>M. intracellulare</i>
2685	ACTCTGATCGTATTGGACACCTAACGTCGAACCCCT-TATC				<i>M. paratuberc.</i>
3385	ACTCTGATCGTATTGGGCACTCTAACCTCGAACCCTGAATC				<i>M. tuberculosis</i>
324	ACTCTGATCGTATTGGGCACTCTAACCTCGAACCCTGAATC				<i>M. bovis</i>
2777	ACTCTGATCGTATTGGGCACTCTAACCTCGAACCCTGAATC				<i>M. phlei</i>
2708	ACTCTGATCGTATTGGGCACTCTAACCTCGAACCCTGAATC				<i>M. leprae</i>
1910					<i>M. gastri</i>
2412	ACTCTGATCGTATTGGACACCTAACGTCGAACCCTGAATC				<i>M. kansasii</i>
4862	ACTCTGATCGTATTGGGCACTCTAACCTCGAACCCTGAATC				<i>M. smegmatis</i>
- - - -					
	2690	2700	2710	2720	
2924	GGTGTCTCAACGGATAAAAGGTACCCCGGGGATAACAG				<i>M. avium</i>
2924	GGTGTCTCAACGGATAAAAGGTACCCCGGGGATAACAG				<i>M. paratuberc.</i>
3625	GGTGTCTCAACGGATAAAAGGTACCCCGGGGATAACAG				<i>M. tuberculosis</i>
3017	GGTGTCTCAACGGATAAAAGGTACCCCGGGGATAACAG				<i>M. phlei</i>
2948	GGTGTCTCAACGGATAAAAGGTACCCCGGGGATAACAG				<i>M. leprae</i>
1910					<i>M. gastri</i>
2652	GGTGTCTCAACGGATAAAAGGTACCCCGGGGATAACAG				<i>M. kansasii</i>
5102	GGTGTCTCAACGGATAAAAGGTACCCCGGGGATAACAG				<i>M. smegmatis</i>
	2730	2740	2750	2760	
2964	GCTGATCTTCCCCAAGAGTCCATATCGACGGGATGGTTTG				<i>M. avium</i>
2964	GCTGATCTTCCCCAAGAGTCCATATCGACGGGATGGTTTG				<i>M. paratuberc.</i>
3665	GCTGATCTTCCCCAAGAGTCCATATCGACGGGATGGTTTG				<i>M. tuberculosis</i>
3057	GCTGATCTTCCCCAAGAGTCCATATCGACGGGATGGTTTG				<i>M. phlei</i>
2988	GCTGATCTTCCCCAAGAGTCCATATCGACGGGATGGTTTG				<i>M. leprae</i>
1910					<i>M. gastri</i>
2692	GCTGATCTTCCCCAAGAGTCCATATCGACGGGATGGTTTG				<i>M. kansasii</i>
5142	GCTGATCTTCCCCAAGAGTCCATATCGACGGGATGGTTTG				<i>M. smegmatis</i>

Figure 4K

27/31

	2770	2780	2790	2800	
3004	GCACCTCGATGTCGGCTCGTCGCATCCTGGGGCTGGAGCA				M. avium
3004	GCACCTCGATGTCGGCTCGTCGCATCCTGGGGCTGGAGCA				M. paratuberc.
3705	GCACCTCGATGTCGGCTCGTCGCATCCTGGGGCTGGAGCA				M. tuberculosis
3097	GCACCTCGATGTCGGCTCGTCGCATCCTGGGGCTGGAGCA				M. phlei
3028	GCACCTCGATGTCGGCTCGTCGCATCCTGGGGCTGGAGCA				M. leprae
1910					M. gastri
2732	GCACCTCGATGTCGGCTCGTCGCATCCTGGGGCTGGAGCA				M. kansasii
5182	GCACCTCGATGTCGGCTCGTCGCATCCTGGGGCTGGAGCA				M. smegmatis
	2810	2820	2830	2840	
3044	GGTCCCAAGGGTTGGGCTGTTTCGCCC-ATTAAAGCGGCAC				M. avium
3044	GGTCCCAAGGGTTGGGCTGTTTCGCCC-ATTAAAGCGGCAC				M. paratuberc.
3745	GGTCCCAAGGGTTGGGCTGTTTCGCCC-ATTAAAGCGGCAC				M. tuberculosis
3137	GGTCCCAAGGGTTGGGCTGTTTCGCCC-ATTAAAGCGGCAC				M. phlei
3068	GGTCCCAAGGGTTGGGCTGTTTCGCCC-ATTAAAGCGGCAC				M. leprae
1910					M. gastri
2772	GGTCCCAAGGGTTGGGCTGTTTCGCCC-ATTAAAGCGGCAC				M. kansasii
5222	GGTCCCAAGGGTTGGGCTGTTTCGCCC-ATTAAAGCGGCAC				M. smegmatis
- - - -					
	3050	3060	3070	3080	
3283	CAAGATCAGGTTT-CTCACCCCTTTTAGAGGGATAAGGCC				M. avium
638	CAAGATCAGGTTT-CTCACCCCTTTTAGAGGGATAAGGCC				M. intracellulare
3283	CAAGATCAGGTTT-CTCACCCCTTTTAGAGGGATAAGGCC				M. paratuberc.
3984	CAAGATCAGGTTT-CTCACCCCTTTTAGAGGGATAAGGCC				M. tuberculosis
570	CAAGATCAGGTTT-CTCACCCCTTTTAGAGGGATAAGGCC				M. bovis
3376	CAAGATCAGGTTT-CTCACCCCTTTTAGAGGGATAAGGCC				M. phlei
3307	CAA				M. leprae
1910					M. gastri
3011	CAAGATCAGGTTT-CTCACCCCTTTTAGAGGGATAAGGCC				M. kansasii
5462	CAAGATCAGGTTT-CTCACCCCTTTTAGAGGGATAAGGCC				M. smegmatis
	3090	3100	3110	3120	
3322	CCCGC-AGACCACGGGATTGATAGGCCAGACCTGGAAGCT				M. avium
677	CCCGC-AGACCACGGGATTGATAGGCCAGACCTGGAAGCT				M. intracellulare
3322	CCCGC-AGACCACGGGATTGATAGGCCAGACCTGGAAGCT				M. paratuberc.
4023	CCCGC-AGACCACGGGATTGATAGGCCAGACCTGGAAGCT				M. tuberculosis
609	CCCGC-AGACCACGGGATTGATAGGCCAGACCTGGAAGCT				M. bovis
3415	CCCGC-AGACCACGGGATTGATAGGCCAGACCTGGAAGCT				M. phlei
3309					M. leprae
1910					M. gastri
3050	CCCGC-AGACCACGGGATTGATAGGCCAGACCTGGAAGCT				M. kansasii
5501	CCCGC-AGACCACGGGATTGATAGGCCAGACCTGGAAGCT				M. smegmatis

Figure 4L

28/31

	130	140	150	160	
107	GAGTAACACGTGGG	CAATCTGCCCTGCACTTC	-GGGATAA		<i>M. avium</i>
59	GAGTAACACGTGGGCAATCTGCCCTGCACTTC	-GGGATAA			<i>M. intracellulare</i>
107	GAGTAACACGTGGGCAATCTGCCCTGCACTTC	-GGGATAA			<i>M. paratuberc.</i>
70	GAGTAACACGTGGGCAATCTGCCCTGCACTTC	-GGGATAA			<i>M. scrofulaceum</i>
70	GAGTAACACGTGGG	TGATCTGCCCTGCACTTC	-GGGATAA		<i>M. tuberculosis</i>
209	GAGTAACACGTGGG	TGATCTGCCCTGCACTTC	-GGGATAA		<i>M. bovis</i>
120	GAGTAACACGTGGG	TAATCTGCCCTGCACTTC	-GGGATAA		<i>M. leprae</i>
69	GAGTAACACGTGGGCAATCTGCCCTGCACACC	-GGGATAA			<i>M. kansasii</i>
70	GAGTAACACGTGGGCAATCTGCCCTGCACACC	-GGGATAA			<i>M. gastri</i>
104	GAGTAACACGTGGG	TAATCTGCCCTGCACATC	-GGGATAA		<i>M. gordonae</i>
64	GAGTAACACGTGGG	CAATCTGCCCTGCACTTC	-GGGATAA		<i>M. marinum</i>

- - - -

	450	460	470	480	
424	AAACCTCTTTTACCATCGACGAAGGTCCGGG	TTTTCTCGG			<i>M. avium</i>
376	AAACCTCTTTTACCATCGACGAAGGTCCGGGTTTTCTCGG				<i>M. intracellulare</i>
424	AAACCTCTTTTACCATCGACGAAGGTCCGGGTTTTCTAGG				<i>M. paratuberc.</i>
387	AAACCTCTTTTACCATCGACGAAGGTCTCA	---CTTTGTGG			<i>M. scrofulaceum</i>
389	AAACCTCTTTTACCATCGACGAAGGTCCGGGTTCTCTCGG				<i>M. tuberculosis</i>
528	AAACCTCTTTTACCATCGACGAAGGTCCGGGTTCTCTCGG				<i>M. bovis</i>
439	AAACCTCTTTTACCATCGACGAAGGTCTGGGA	ATTCTCGG			<i>M. leprae</i>
386	AAACCTCTTTTACCATCGACGAAGGTCCGGGTTCTCTCGG				<i>M. kansasii</i>
387	AAACCTCTTTTACCATCGACGAAGGTCCGGGTTCTCTCGG				<i>M. gastri</i>
420	AAACCTCTTTTACCATCGACGAAGGTCCGGGTTTTCTCGG				<i>M. gordonae</i>
381	AAACCTCTTTTACCATCGACGAAGGTCTCGGGTTTTCTCGG				<i>M. marinum</i>

	490	500	510	520	
429	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. tuberculosis</i>
568	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. bovis</i>
464	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. avium</i>
416	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. intracellulare</i>
464	ATTGACGGTAGGTGGAGAAGAAGCAC	-----	ACTACGTG		<i>M. paratuberc.</i>
424	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. scrofulaceum</i>
479	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. leprae</i>
426	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. kansasii</i>
427	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. gastri</i>
460	GCTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. gordonae</i>
421	ATTGACGGTAGGTGGAGAAGAAGCACCGGCCAACTACGTG				<i>M. marinum</i>

Figure 5A

29/31

	1130	1140	1150	1160	
1104	TCTCATGTTGCCAGC	GGGTAATGC	GGGGACTCGTGAGAG		M.avium
1056	TCTCATGTTGCCAGC	GGGTAATGCCGGGGACTCGTGAGAG			M.intracellulare
1098	TCTCATGTTGCCAGC	GGGTAATGC	GGGGACTCGTGAGAG		M.paratuberc.
1064	TCTCATGTTGCCAGC	GGGTAATGCCGGGGACTCGTGAGAG			M.scrofulaceum
1069	TCTCATGTTGCCAGC	ACGTAATG	GTGGGGACTCGTGAGAG		M.tuberculosis
1208	TCTCATGTTGCCAGC	ACGTAATG	GTGGGGACTCGTGAGAG		M.bovis
1119	TCTCATGTTGCCAGC	ACGTAATG	GTGGGGACTCGTGAGAG		M.leprae
1066	TCTCATGTTGCCAGC	GGGTAATGCCGGGGACTCGTGAGAG			M.kansasii
1067	TCTCATGTTGCCAGC	GGGTAATGCCGGGGACTCGTGAGAG			M.gastri
1100	TCTCATGTTGCCAGC	GGGTAATGCCGGGGACTCGTGAGAG			M.gordonae
1061	TCTCATGTTGCCAGC	ACGTAATG	GTGGGGACTCGTGAGAG		M.marinum

- - - -

	1290	1300	1310	1320	
1264	CGAATCCTTTTAAAGCCGGACTCAGTTCGGAT	TGGGGTCT			M.avium
1216	CGAATCCTTTTAAAGCCGG	CTCAGTTCGGATTGGGGTCT			M.intracellulare
1258	CGAATCCTTTTAAAGCCGGACTCAGTTCGGATTGGGGTCT				M.paratuberc.
1224	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGAT	CGGGTCT			M.scrofulaceum
1229	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGAT	CGGGTCT			M.tuberculosis
1368	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGAT	CGGGTCT			M.bovis
1279	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGAT	CGGGTCT			M.leprae
1226	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGAT	CGGGTCT			M.kansasii
1227	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGAT	CGGGTCT			M.gastri
1260	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGAT	CGGGTCT			M.gordonae
1221	CGAATCCTTTTAAAGCCGGTCTCAGTTCGGAT	CGGGTCT			M.marinum

	1330	1340	1350	1360	
1304	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.avium
1256	GCAACTCGACCCCATGAAGTCGGAGTCGCTAGTAATCGCA				M.intracellulare
1298	GCAACTCGACCCCATGAAGTCGGAGTCGCTAGTAATCGCA				M.paratuberc.
1264	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.scrofulaceum
1268	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.tuberculosis
1407	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.bovis
1319	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.leprae
1266	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.kansasii
1267	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.gastri
1300	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.gordonae
1260	GCAACTCGACCCCGTGAAGTCGGAGTCGCTAGTAATCGCA				M.marinum

Figure 5B

30/31

2550 | TTACGGCGGCAGGACGAAAGACCCCGGACCTTCACTA 2589 |
2568 2569 |
M. avium 23S:

Figure 6

31/31




Figure 7

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 97/00425

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12Q1/68 C07K14/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 547 842 A (HOGAN JAMES ET AL) 20 August 1996 cited in the application see the whole document ---	1-36
Y	WO 96 17956 A (GENE POOL INC ; WEININGER SUSAN (US); WEININGER ARTHUR M (US)) 13 June 1996 see the whole document ---	1-36
Y	WO 95 32305 A (DAKO AS) 30 November 1995 see the whole document ---	1-36
A	EP 0 572 120 A (GEN PROBE INC) 1 December 1993 cited in the application see the whole document ---	
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

20 January 1998

Date of mailing of the international search report

30/01/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Knehr, M

INTERNATIONAL SEARCH REPORT

Inter. Appl. Application No
PCT/DK 97/00425

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	WO 96 36734 A (ABBOTT LAB) 21 November 1996 see the whole document -----	1-36

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. Appl. No.

PCT/DK 97/00425

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5547842 A	20-08-96	US 5541308 A	30-07-96
		US 5595874 A	21-01-97
		US 5593841 A	14-01-97
		US 5683876 A	04-11-97
		US 5677127 A	14-10-97
		US 5677128 A	14-10-97
		US 5677129 A	14-10-97
		US 5693468 A	02-12-97
		US 5691149 A	25-11-97
		US 5693469 A	02-12-97
		US 5679520 A	21-10-97
		US 5674684 A	07-10-97
		DK 413788 A	23-09-88
		AU 616646 B	07-11-91
		AU 1041988 A	16-06-88
		EP 0272009 A	22-06-88
		JP 1503356 T	16-11-89
		KR 9511719 B	09-10-95
		WO 8803957 A	02-06-88
WO 9617956 A	13-06-96	AU 4418996 A	26-06-96
		CA 2206127 A	13-06-96
		EP 0796344 A	24-09-97
		NO 972611 A	11-08-97
WO 9532305 A	30-11-95	AU 2522095 A	18-12-95
		EP 0760008 A	05-03-97
EP 0572120 A	01-12-93	AU 4114793 A	29-11-93
		JP 7506723 T	27-07-95
		WO 9322330 A	11-11-93
WO 9636734 A	21-11-96	NONE	